

Part I

Changing Risk Behaviors and Addressing Environmental Challenges

Chapter 1

Tobacco

Reducing Tobacco Use Initiation

RECOMMENDED INTERVENTIONS

- Increasing the Unit Price for Tobacco Products* 10
- Mass Media Education Campaigns When Combined with Other Interventions* 12
- Restricting Minors' Access to Tobacco Products: Community Mobilization When Combined with Additional Interventions (Stronger Local Laws Directed at Retailers, Active Enforcement of Retailer Sales Laws, Retailer Education with Reinforcement)* 17

INSUFFICIENT EVIDENCE TO DETERMINE EFFECTIVENESS OF THE INTERVENTION*

- Restricting Minors' Access to Tobacco Products: Sales Laws Directed at Tobacco Retailers to Reduce Illegal Sales to Minors When Implemented Alone* 20
- Laws Directed at Minors' Purchase, Possession, or Use of Tobacco Products When Implemented Alone* 21
- Active Enforcement of Sales Laws Directed at Retailers When Implemented Alone* 22
- Retailer Education with Reinforcement and Information on Health Consequences When Implemented Alone* 23
- Retailer Education without Reinforcement When Implemented Alone* 24
- Community Education about Minors' Access to Tobacco Products When Implemented Alone* 25

Increasing Tobacco Use Cessation

RECOMMENDED INTERVENTIONS

- Increasing the Unit Price for Tobacco Products* 27
- Mass Media Education Campaigns Combined with Other Interventions* 29
- Healthcare Provider Reminder Systems* 35
- Healthcare Provider Reminder Systems with Provider Education, with or without Client Education* 37
- Reducing Client Out-of-Pocket Costs for Effective Cessation Therapies* 42
- Multicomponent Interventions That Include Client Telephone Support* 44

*Insufficient evidence means that we were not able to determine whether or not the intervention works.

The Task Force approved the recommendations in this chapter in 1999–2000. The research on which the findings are based was conducted between 1976 and May 2000. This information—with the exception of reviews of approaches to reduce tobacco use through interventions to restrict minors' access to tobacco products—has been previously published in the *American Journal of Preventive Medicine* [2001; 20(2S): 10–15 and 16–66] and the *MMWR Recommendations and Reports* series [2000; 49(No. RR-12):1–11]. The information on restricting minors' access to tobacco products is being prepared for publication.

INSUFFICIENT EVIDENCE TO DETERMINE EFFECTIVENESS OF THE INTERVENTION*

<i>Mass Media Education—Cessation Series</i>	32
<i>Mass Media Education—Cessation Contests</i>	34
<i>Healthcare Provider Education Alone</i>	39
<i>Healthcare Provider Feedback and Assessment</i>	41

Reducing Exposure To Environmental Tobacco Smoke

RECOMMENDED INTERVENTIONS

<i>Smoking Bans and Restrictions</i>	48
--------------------------------------	----

INSUFFICIENT EVIDENCE TO DETERMINE EFFECTIVENESS OF THE INTERVENTION*

<i>Community Education to Reduce Exposure to Environmental Tobacco Smoke in the Home</i>	51
--	----

Tobacco use is the leading cause of preventable illness and death in the United States.¹ Recognized as a cause of multiple cancers, heart disease, stroke, complications of pregnancy, and chronic obstructive pulmonary disease,² tobacco use is responsible for 430,000 deaths per year among adults, and direct medical costs are in the range of \$50–\$73 billion per year.³

Despite nearly four decades of policies, regulations, educational efforts, and increasing information on the negative health effects of tobacco use and the positive health benefits of cessation, tobacco use remains unacceptably high. In 2002, an estimated 45.8 million adults in the United States (nearly one in four adults) smoked.⁴ Tobacco use and disparities in tobacco-related morbidity and mortality vary by region, level of education, socioeconomic status, race, gender, and ethnicity.^{2,5} Individuals below the poverty line, for example, are more likely to smoke than individuals at or above the poverty line (32.9% compared with 22.2%).⁴ By level of education, adults who had earned a General Educational Development (GED) diploma had the highest prevalence (42.3%) of smoking; people with masters, professional, and doctoral degrees had the lowest prevalence (7.2%).⁴ The prevalence of smoking among American Indians and Alaska Natives (40.8%) is higher than that of other racial and ethnic groups.⁴

Regular tobacco use results in true drug dependence in most users, making attempts to quit difficult and relapses common.⁶ Many users make multiple attempts to quit.⁷ In 2002, an estimated 15.4 million current smokers (41.2%) had stopped smoking for at least one day during the preceding 12 months because they were trying to stop smoking entirely.⁴ Although cessation significantly reduces the immediate and subsequent risks of tobacco-related morbidity and mortality,^{8,9} most tobacco users do not receive assistance in quitting.⁶

Preventing the acquisition of this costly, chronic dependence is clearly de-

sirable. However, tobacco use initiation and the transition from experimentation to addiction are not easy to prevent because they occur primarily in adolescence, when individuals are more susceptible to influences from family, friends, peers, society, and the tobacco industry that encourage tobacco use.¹⁰ Smoking among 8th, 10th, and 12th graders increased dramatically from 1991 to 1997 and then declined just as dramatically from 1997 to 2003. For example, prevalence of current smoking among high school seniors was 28.3% in 1991, increased to 36.5% by 1997, and declined to 24.4% by 2003.¹¹ The high rates of smoking prevalence among young adults aged 18–24 years (28.5% in 2002), in addition to reflecting those who started smoking in high school, may also indicate an increase in tobacco use initiation among this group.⁴

Exposure to environmental tobacco smoke (ETS) is a recognized cause of heart disease and accounts for an estimated 3000 lung cancer deaths per year among adults.¹² Among infants and children, exposure to ETS causes middle ear infections and effusions, exacerbates 400,000–1,000,000 cases of asthma annually, and causes 150,000–300,000 cases of lower respiratory tract infections each year.¹³ The health effects of exposure to ETS have prompted the increasing implementation of public and private policies that restrict smoking.² The median level of cotinine (a marker for exposure to tobacco smoke) among nonsmokers declined 70% from 1988–1991 to 1999–2000, suggesting a dramatic reduction in exposure of the general U.S. population to ETS over this time period. However, substantial gender, racial, and age disparities in exposure are evident, with children, men, and African Americans having higher exposure levels.¹⁴ Exposure to ETS continues to occur in workplaces and public areas without smoking bans or effective restrictions and in households in which smoking indoors is allowed.

OBJECTIVES AND RECOMMENDATIONS FROM OTHER ADVISORY GROUPS

The interventions reviewed in this chapter can be useful in reaching many of the tobacco control objectives in *Healthy People 2010*¹⁵ (Table 1–1). In the field of tobacco control, many groups have published useful guidelines. To help summarize this information, we published an article in 2001¹⁶ comparing the *Community Guide* population-based recommendations with reviews and recommendations produced by other groups, including the Surgeon General's office, the United States Preventive Services Task Force (which publishes the *Guide to Clinical Preventive Services*¹⁷), and the Cochrane Collaboration. Another good source for readers interested in information on various clinical guidelines is *Treating Tobacco Use and Dependence: Clinical Practice Guideline*⁶; some information from that report is also summarized in our article.

Table 1–1. Selected *Healthy People 2010*¹⁵ Objectives Related to Tobacco Use and Environmental Tobacco Smoke Exposure

<i>Objective</i>	<i>Population</i>	<i>Baseline</i>	<i>2010 Objective</i>
<i>Tobacco Use in Population Groups</i>			
Reduce cigarette smoking by adults (Objective 27–1a)	Adults	24% (1998)	12%
Reduce use of tobacco products by adolescents (in the past month)(27–2a)	Adolescents	40% (1999)	21%
Reduce the initiation of tobacco use among children and adolescents (27–3)	Children/adolescents	Developmental	
Increase (delay) the average age of first tobacco use by adolescents (aged 12–17 years) (27–4a)	Adolescents	12 years old (1997)	14 years old
<i>Cessation and Treatment</i>			
Increase the percentage of smokers stopping smoking for a day or longer (27–5)	Adults	41% (1998 ^a)	75%
Among females aged 18–49 years, increase smoking cessation in pregnant women in the first trimester of pregnancy (27–6)	Pregnant women	14% (1998)	30%
Among adolescents in grades 9–12, increase the percentage of ever-daily smokers who try to quit (27–7)	Adolescents	76% (1999)	84%
Increase in insurance coverage of evidence-based treatment for nicotine dependency:			
• in managed care organizations: (27–8a)	All	75% (1997–98)	100%
• in Medicaid programs in states and the District of Columbia (27–8b)		24 states (1998)	51 states
<i>Exposure to Secondhand Smoke (ETS)</i>			
Among children aged ≤6 years, reduce the proportion of children who live in homes in which someone smokes inside the house ≥4 days per week (27–9)	Young children	27% (1994)	10%
Reduce the proportion of nonsmokers aged ≥4 years with a serum cotinine level >0.10 ng/mL (27–10)	Nonsmokers	65% (1988–94 ^a)	45%

Table 1-1. Continued

<i>Objective</i>	<i>Population</i>	<i>Baseline</i>	<i>2010 Objective</i>
<i>Exposure to Secondhand Smoke (ETS) (continued)</i>			
Increase the proportion of smoke-free and tobacco-free middle, junior high, and senior high schools (27-11)	Schools	37% (1994)	100%
Increase the proportion of worksites (with ≥ 50 workers) with formal smoking policies that prohibit smoking or limit it to separately ventilated areas (27-12)	Employees	79% (1998-99)	100%
Increase the number of states plus the District of Columbia with laws on smoke-free air:		(All 1998)	(For all)
• in private workplaces (27-13a)	All	1 state	51 states
• in public workplaces (27-13b)		13 states	
• in restaurants (27-13c)		3 states	
• on public transportation (27-13d)		16 states	
• in day-care centers (27-13e)		22 states	
• in retail stores (27-13f)		4 states	
<i>Social and Environmental Changes</i>			
Increase the number of states (including the District of Columbia) with a $\leq 5\%$ sales rate of tobacco products to minors (27-14a)	Minors	0 states (1998)	51 states
Increase the number of states (including the District of Columbia) that can suspend or revoke state licenses for violation of laws prohibiting the sale of tobacco to minors (27-15)	Minors	34 states (1998)	51 states
Eliminate tobacco advertising and promotions that influence adolescents and young adults (27-16)	Adolescents/ young adults	Developmental	
Increase the proportion of adolescents who disapprove of smoking:		(All 1998)	
• among 8th graders (27-17a)	Adolescents	80%	95% for all

continued next page

Table 1–1. *Continued*

<i>Objective</i>	<i>Population</i>	<i>Baseline</i>	<i>2010 Objective</i>
<i>Social and Environmental Changes (continued)</i>			
• among 10th graders (27–17b)		75 %	
• among 12th graders (27–17c)		69 %	
Increase the number of tribes, territories, and states and the District of Columbia with comprehensive, evidence-based tobacco control programs (27–18)	All	Developmental	
Reduce the number of states that have laws preempting stronger tobacco control (in the areas of clean indoor air, minors' access laws, or marketing) (27–19)	All	30 states (1998)	0 states
Reduce the toxicity of tobacco products by establishing a regulatory structure to monitor toxicity (27–20)	All	Developmental	
• Increase the combined federal and average state tax (per pack) on cigarettes (27–21a)	All	\$0.63 (1998)	\$2
• Increase the combined federal and average state tax on spit tobacco (27–21b)		Developmental	

^aAge adjusted for the year 2000 standard population.

METHODS

Methods used for the reviews are summarized in Chapter 10. Specific methods used in the systematic reviews of tobacco use prevention and control have been described elsewhere^{16,18–20} and are available at www.thecommunityguide.org/tobacco. The logic framework depicting the conceptual approach used in these reviews is presented in Figure 1–1.

ECONOMIC EFFICIENCY

A systematic review of available economic evaluations was conducted for each recommended intervention, and a summary of each review is presented with the related intervention. The methods used to conduct these economics reviews are summarized in Chapter 11.




Figure 1–1. Logic framework illustrating the conceptual approach used in systematic reviews of tobacco use. (ETS = environmental tobacco smoke.) (Reprinted from Am J Prev Med, Vol. 20, No. 2S, Hopkins DP et al., Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke, p. 18, Copyright 2001, with permission from American Journal of Preventive Medicine.)

RECOMMENDATIONS AND FINDINGS

This section presents a summary of the findings of the systematic reviews conducted to determine the effectiveness of the selected interventions in this topic area. We evaluated interventions to address three goals essential to tobacco prevention and control efforts: decreasing the number of people who start using tobacco products (tobacco use initiation), increasing the number of tobacco users who quit (tobacco use cessation), and reducing exposures among nonsmokers to environmental tobacco smoke (ETS).

Reducing Tobacco Use Initiation

The interventions in our reviews are designed to change knowledge, attitudes, and tobacco use among children, adolescents, and young adults. Most adults who use tobacco products began in adolescence, and nicotine addiction develops during the first few years of use.²¹ Children and adolescents may perceive tobacco use to be a normal peer and adult behavior, and can often act on this belief because tobacco products are readily available and accessible.¹⁰ Preventing or delaying experimentation with tobacco or preventing the transition from experimentation to regular use are the major goals of the interventions we reviewed: increasing the unit price for tobacco products,

mass media campaigns when combined with additional efforts, and interventions to restrict minors' access to tobacco products.

***Increasing the Unit Price for Tobacco Products:
Recommended (Strong Evidence of Effectiveness)***

Interventions to increase the unit price for tobacco products primarily include legislation at the municipal, state, or federal level that raises the excise tax on tobacco products. In several states, excise tax increases have resulted from successful state ballot initiatives. Although other factors affect tobacco product pricing, excise tax increases have historically resulted in an equivalent or larger increase in tobacco product prices.²²

Excise taxes on tobacco products increase the overall product cost and therefore make the use of tobacco products less attractive to young people with limited income and a variety of ways to spend their money.

Effectiveness

- Higher prices for tobacco products are associated with reduced consumption of tobacco products (a 10% price increase results in approximately a 4% decrease).
- Higher prices for tobacco products are associated with lower levels of tobacco use by adolescents and young adults who use tobacco (a 10% price increase results in approximately a 2% decrease in the amount smoked).

Applicability

- These findings should be applicable to most adolescents and young adults in the United States.

Other Effects

- Price increases also reduce tobacco use among adults.

Barriers

- Increasing the excise tax on tobacco products requires passage of legislation or a statewide referendum.

The findings of our systematic review are based on eight studies that evaluated the effectiveness of increasing the price of tobacco products on changing the tobacco use behaviors of adolescents or young adults.²³⁻³⁰ Three additional articles provided information on a study already included in the review.³¹⁻³³ All studies were conducted in the United States. Studies examined the effect of product price on tobacco use by adolescents (13-18 years old) and by young adults (18-24 years or under 25 years old).

All eight studies used econometric methods to analyze surveys of students

or young adults. Price elasticity of demand (the percentage change in consumption that results from a 1% change in price) estimates were calculated by combining information on local tobacco product prices, price changes, or differences over the period of study with responses to surveys on tobacco use and consumption. The price elasticity of demand estimates in these studies included participation (i.e., tobacco use prevalence), number of cigarettes smoked per day among those who smoke, and overall estimates of tobacco consumption (which includes both changes in participation and amount smoked by those who continue to smoke). All of the studies attempted to control for concurrent tobacco prevention and control efforts, including differences in smoking restrictions, youth tobacco access laws, school tobacco education programs, and exposure to anti-tobacco media. Five studies evaluated the effect of price on tobacco use for study periods that included the 1990s, and three studies reported the effect of price on tobacco use for periods before 1990.

Price elasticity of demand estimates from seven studies demonstrated that higher tobacco product prices are associated with lower levels of tobacco use by adolescents and young adults. Based on the median estimate of price elasticity from the studies in this review, for example, tobacco use prevalence among adolescents (13–18 years old) would decrease 3.7% with a 10% increase in product price (range, “no statistically significant effect” to 11.9%). The effects are at least as strong for young adults (18–24 years old). For estimates of amount smoked, the median estimate from six studies suggests that a 10% increase in product price would result in a 2.3% decrease in the amount of tobacco used by adolescents who smoke.

These findings show that increasing the unit price for tobacco products is effective in reducing both tobacco use prevalence and tobacco consumption among adolescents and young adults.

These results should be applicable to most adolescents and young adults in the United States. All studies were conducted in the United States, and most of the studies used national datasets to compare tobacco use and product price across jurisdictions (e.g., states). The study samples are representative of populations of adolescents and young adults. Studies evaluated the effect of product price on use and consumption of both cigarettes and smokeless tobacco products.

Overall, the studies showed that tobacco product price increases had a greater effect among males than among females; price affected tobacco use and consumption among whites, African Americans, and Hispanics; and African-American adolescents and young adults were more responsive to differences in product price than were white adolescents and young adults, respectively.

Increasing the price of tobacco products also reduces tobacco use among adults (for additional information see the review of Increasing the Unit Price for Tobacco Products in the next section, Increasing Tobacco Use Cessation). We found no other additional positive or negative effects of tobacco product price increases on adolescent tobacco use initiation.

Econometric analyses were included in the studies qualifying for the review of effectiveness. A separate economic evaluation was not conducted for this intervention review.

Passage of legislation or a statewide referendum is required for an excise tax increase on tobacco products and may, therefore, present a significant barrier to implementation. Political opposition has historically been well organized and funded at both the federal and state levels. Reports on both successful and unsuccessful state initiatives that proposed an increase on tobacco product excise taxes have been published.³⁴⁻³⁶

In conclusion, the Task Force recommends increasing the unit price for tobacco on the basis of strong evidence of effectiveness in reducing tobacco use initiation among adolescents and young adults. Raising the unit price for tobacco products (through increases in the excise tax) is effective in reducing both the number of adolescents and young adults who use tobacco products and the amount of tobacco they use. Increasing the unit price for tobacco products also reduces tobacco use among adults. The need for passage by a state legislature or a referendum can present a barrier to increasing excise taxes.

Mass Media Education Campaigns When Combined with Other Interventions: Recommended (Strong Evidence of Effectiveness)

These campaigns use mass media for an extended duration (months to years) to disseminate brief, recurring messages with the goal of providing information that will motivate people, primarily children and adolescents, to remain tobacco-free. The messages, developed through formative research, are disseminated through paid broadcast time and print space, donated time and space (as public service announcements), or a combination of the two. Mass media campaigns can be combined or coordinated with additional interventions, such as increases in tobacco product excise taxes, school-based education, and other community-wide educational activities.

Mass media techniques primarily include broadcast messages on television and radio, although other formats such as billboards, print, and movies have been used. Campaigns can focus on messages targeting children and adolescents or can include such messages as part of an overall anti-tobacco effort (e.g., including messages targeting tobacco users to increase cessation

and messages about reducing exposure to secondhand tobacco smoke). The content of mass media campaigns designed to educate and motivate children and adolescents to remain tobacco-free can vary, but a recent review identified two primary strategies: agenda setting and demand reduction education.³⁷ Agenda-setting messages increase awareness of strategies used by the tobacco industry to promote tobacco use, and attempt to facilitate changes in both tobacco use behaviors and public tobacco policies. Demand reduction education messages provide information and support to young people to help them decide to remain tobacco-free.

Effectiveness

- Mass media education campaigns, combined with other interventions, are effective in decreasing the number of young people who use tobacco by approximately 2.4 percentage points.
- Effectiveness was increased in campaigns lasting two years or longer.

Applicability

- These findings should be applicable to most adolescents in the United States.

Other Effects

- Mass media education campaigns can also include messages that contribute to reductions in tobacco use among adults.

The findings of our systematic review are based on 12 studies that evaluated the effectiveness of mass media campaigns in reducing tobacco use among adolescents.³⁸⁻⁴⁹ Two additional studies were identified but did not meet our quality criteria and were excluded from the review.^{50,51} Another 28 papers provided information on studies already included in the review.⁵²⁻⁷⁹ In nine studies, the mass media efforts focused on youth; in three other studies, youth-targeted messages were part of a larger anti-tobacco campaign. Only one study used mass media alone (through a variety of outlets). In 11 studies, the mass media campaign was conducted in coordination or concurrently with contests, school-based education programs, or community education programs. Two studies were conducted in settings where excise taxes on tobacco products had been increased. Two of the mass media campaigns were less than two weeks long, three were less than two years long, and seven campaigns were more than two years long.

The qualifying studies provided 12 diverse measures of tobacco use and generally showed reductions among youth (students) in the intervention communities relative to youth in the comparison communities. For example, five studies reported absolute differences in self-reported tobacco use and observed a median reduction of 2.4 percentage points (range, 0.02 increase to

9.5 decrease). In six studies, comparisons in self-reported tobacco use were expressed as odds ratios. Two of these studies reported no intervention effect on tobacco use, whereas four studies observed lower rates among people exposed to the intervention (median odds ratio of 0.60, range 0.49 to 0.74).

All seven studies evaluating campaigns of two or more years' duration observed reductions in tobacco use among youth in favor of the intervention communities. In studies reporting absolute percentage differences the median reduction in tobacco use was 8.0 percentage points (range, 2.4 to 11), and in studies calculating odds ratios the median was 0.74 (range, 0.49 to 0.74).

We could not assess the effects of the separate components of these combined intervention efforts.

These findings show that mass media education campaigns, when combined with other intervention activities, are effective in decreasing tobacco use among youth.

These results should be applicable to most adolescents in the United States. Statewide campaigns were conducted in Florida, Massachusetts, and Minnesota. Community and regional interventions were conducted in the Southeast, Northeast, and Midwest, and in Montana and Southern California. Campaigns began in the 1990s, the 1980s, and the 1970s. Both representative samples of adolescents in the general population and representative or selected samples of schools were used. Student populations were recruited or surveyed before grade 6, in grades 6–7, or after grade 7. The two most recent studies showed a greater benefit among younger adolescents. Evaluation of the Florida campaign provides evidence of effectiveness of mass media campaigns among girls, boys, whites, African Americans, and Hispanics.

Mass media campaigns, when combined with additional interventions, can also reduce tobacco use among adults (for additional information see Mass Media Education Campaigns in the next section, Increasing Tobacco Use Cessation), although the message content, broadcast times, and settings that reduce youth tobacco use may not be effective in reducing adult tobacco use. No other positive or negative effects were identified.

The findings of our systematic review of economic evaluations are based on one four-year economic study conducted in Montana, New York, and Vermont.⁷¹ This cost-effectiveness analysis compared the effect on students of a mass media campaign combined with a school smoking prevention program to the effect of a school smoking prevention program alone. Students (from grades 5–7 through grades 8–10) were followed for two years after the intervention. Costs included personnel, travel, data entry, message research and development, and television and radio advertising. At follow-up, smoking prevalence among students exposed to the media intervention plus the school

smoking prevention program was 20.4%, compared with 25.9% among students exposed only to the school program. Adjusted program cost per smoker averted was \$6069, and the adjusted program cost per quality-adjusted life year (QALY) was \$333.

The main barrier to implementation of mass media campaigns is the cost of purchasing advertising time. Costs of developing and test marketing messages can be offset by cooperation between tobacco control programs.

In conclusion, the Task Force recommends mass media campaigns, when combined with additional interventions, on the basis of strong evidence of effectiveness in reducing tobacco use among adolescents. These interventions were effective in decreasing the number of young people who use tobacco products. The findings of this review should be applicable to most adolescents in the United States. Mass media campaigns can produce the additional benefit of a reduction in adult tobacco use.

Restricting Minors' Access to Tobacco Products: Interventions to Reduce Tobacco Use

Access to tobacco products by minors (people under 18 years of age in most states) contributes to the initiation and regular use of tobacco by children and adolescents.^{2,10} Retailers who sell tobacco products to minors (including vending machines in accessible settings) constitute one avenue of access for minors.^{2,10,80} Social sources provide another route of access for many, but not all, minors. Although social sources include adults (parents, family, friends) who may purchase tobacco products legally,⁸¹⁻⁸³ illegal retailer sales to minors provide tobacco products for distribution to other minors, contributing to social access.⁸¹

Minors obtain tobacco from commercial sources through face-to-face purchases from retailers (from self-service displays or requests for products held behind the counter), purchases from vending machines, purchases through the mail or over the Internet, access to free product samples, and theft from retail sources. Overall, minors' access reflects the availability of tobacco products within the community, the willingness of retailers to sell them, and the efforts of minors to obtain them. Interventions to reduce access attempt to modify or to change one or more of these factors.

This section includes a review of a variety of interventions to restrict and reduce the supply of tobacco products that minors can obtain from commercial sources. Some of the intervention combinations reviewed included components intended, in whole or in part, to reduce the demand for tobacco products by minors through efforts to educate and mobilize the community

and to change social norms about the acceptability of tobacco use.⁸⁴ In these efforts, minors' access to commercial sources of tobacco provided a focus for the community to address the problem of tobacco use among youth.

In conducting this review of interventions to reduce minors' access to tobacco products, we considered as evidence of effectiveness (1) measurements of differences or changes in self-reported tobacco use in the study population and (2) measurements of differences or changes in self-reported purchases of tobacco products or use of tobacco obtained from commercial sources. Because it is not clear how the compliance of any particular retailer contributes to the overall availability of tobacco to youth, differences or changes in retailer sales of tobacco products to minors were evaluated and summarized but did not by themselves provide the basis for our assessments of effectiveness.

In this section, we review studies that combined two or more interventions in a coordinated effort to restrict minors' access to tobacco products. We identified a total of 28 multicomponent studies. The 13 studies (with 14 arms) that qualified for our assessment of effectiveness evaluated a total of 10 different combinations of interventions. However, only five of these studies conducted measurements of differences or changes in tobacco use among youth, the focus of our assessment of effectiveness. The remaining studies measured differences or changes in retailer sales rates on "youth test purchase attempts" (in which recruited, trained, and supervised minors make standardized attempts to purchase tobacco products from retailers and vending machines). Overall, the body of evidence included evaluations of effectiveness for a variety of overlapping combinations of interventions. The information presented here summarizes only one of our reviews on effectiveness across the qualifying multicomponent studies.

Community Mobilization When Combined with Additional Interventions: Recommended (Sufficient Evidence of Effectiveness)

These interventions are implemented community-wide to focus public attention on the issue of youth access to tobacco products and to generate and mobilize community support for additional efforts to reduce that access. The community mobilization efforts evaluated here either fostered or were coordinated with additional interventions, such as stronger restrictions on retailer sales of tobacco products; restrictions directed at youth purchase, possession, or use; active enforcement of tobacco sales laws; and retailer education interventions (with or without reinforcement).

Educational components of the interventions included community-wide assessments of compliance by tobacco retailers—with dissemination of the results through mass media events and news coverage—and presentations to

civic groups and local governments. Community and school meetings and activities, as well as direct contact with local governments through testimony, petitions, letters, and phone calls, also occurred.

We included studies in the review regardless of the order in which components were added. For example, in some studies, community mobilization was the initial intervention and contributed to the adoption of additional interventions. In other studies, community mobilization followed other interventions (such as stronger laws directed at retailers) in efforts to generate and maintain community support for reducing minors' access to tobacco products.

Effectiveness

- Community mobilization, when combined with other interventions, was effective in reducing tobacco use among youth (students) by approximately 5.8 percentage points.
- The intervention reduced the sale of tobacco products by retailers to youth making test purchase attempts by approximately 34 percentage points.

Applicability

- These findings should be applicable to most communities in the United States where not precluded by preemption legislation.

Barriers to Implementation

- The adoption or existence of a law at the state level that supersedes or precludes stronger local laws (preemption) is a significant barrier to the effective combination of community mobilization and coordinated interventions.

We identified a total of nine qualifying studies (10 intervention arms) that provided evidence on the effectiveness of community mobilization when combined with additional interventions.⁸⁵⁻⁹³ Seven additional papers provided information on studies already included in our review.⁹⁴⁻¹⁰⁰ Seven studies did not meet our quality criteria and were excluded from the review.¹⁰¹⁻¹⁰⁷

As noted above, most of the qualifying studies did not attempt to measure differences or changes in tobacco use among youth, the focus of our assessment of effectiveness. The nine qualifying studies provided 10 measurements of changes in the percentage of retailers willing to sell tobacco products to youth in test purchase attempts. Overall, the studies observed a 33.5 percentage point reduction in retailers willing to sell tobacco to youth (range, 4.5 to 68). In the intervention communities at follow-up, the median sales rate among retailers on youth test purchase attempts was 27.5% (range, 0% to 65%).

The findings of this systematic review are based on the subset of five multi-component intervention studies that measured differences or changes in tobacco use among youth in the study communities.^{86,87,90,91,108} Four of these studies evaluated a combined intervention including community mobiliza-

tion.^{86,87,90,91} In these studies, the interventions coordinated with community mobilization included retailer education with reinforcement,^{86,87,91} stronger local ordinances directed at retailers,^{90,91} active enforcement of retailer sales laws,^{90,91} local ordinances directed at youth tobacco purchase, possession, or use,^{90,91} and school-based education.⁸⁷

All four studies evaluating interventions combined with community mobilization observed decreases in self-reported tobacco use by students, with a median decrease of 5.8 percentage points (range, 3.8 to 11) over follow-up periods of 24 – 48 months. In addition, all four studies observed both reductions in the sale of tobacco products on youth test purchase attempts (median – 44 percentage points) and low post-intervention sales rates by retailers on youth test purchase attempts (median 7.9%; range, 0% to 24%) in the intervention communities.

The fifth qualifying study that measured youth tobacco use as an intervention outcome evaluated the combination of sustained, active enforcement of sales laws directed at retailers following a brief retailer education effort (notification of existing sales laws).¹⁰⁸ Despite a significant reduction of 30 percentage points in retailer sale of tobacco products on youth test purchase attempts (the post-intervention sales rate was 18% in the intervention community), self-reported tobacco use by surveyed high school students was not significantly different at two-year follow-up.

Overall, these results indicate that the combination of community mobilization and additional activities is effective in reducing youth access to tobacco.

These results should be applicable to some communities in the United States (barriers are described below). The multicomponent interventions evaluated in this body of evidence were implemented in a variety of settings and populations, including urban, suburban, and rural communities in the United States and Australia. Interventions were implemented in U.S. communities that included predominantly African-American, Hispanic, or white populations.

We found no information on other positive or negative effects of community mobilization. One report observed that youth who took part in test purchase attempts had not increased smoking or intentions to smoke at a two-year follow-up.¹⁰⁹

The findings of our systematic review of economic evaluations are based on one study of a component of community mobilization.¹¹⁰ This one-year study modeled the cost effectiveness of active enforcement of tobacco sales to minors on a national level. The intervention included employing minors to attempt tobacco purchases, licensing tobacco vendors, and civil penalties for vendors who illegally sold tobacco products to minors. Program costs in-

cluded personnel, salary, and benefits for minors and for adult inspectors; liability insurance; money to purchase tobacco; transportation; and overhead (analyses were based on enforcement costs of \$50, \$150, \$250, and \$350, where marginal expense is lowest at the community level and highest at the federal level). Primary outcome measures consisted of four levels of reduction in youth tobacco use ranging from 5% to 50%, with subsequent cost-effectiveness ratios ranging from \$44 to \$3100 per year of life saved.

Barriers to implementation are well described in the published literature. Preemption, the adoption or existence of a law at the state level that supersedes or precludes stronger local laws, is the subject of a *Healthy People 2010*¹⁵ objective (see Table 1-1). Preemption is a significant barrier to the effective combination of community mobilization and coordinated interventions, such as laws directed at retailers and the conduct of active enforcement.^{2,90,106,111,112} Preemption statutes in many states hinder or prevent implementation of the effective combinations of interventions described in this review.^{2,106} In addition, some preemption laws hinder or obstruct compliance checks of retailers, making it more difficult for communities to recognize and address minors' access to commercial sources of tobacco products.

Opposition by retailers, retail associations, and the tobacco industry can be a significant barrier to the implementation and conduct of sustained, active enforcement of sales laws.^{2,90} Lack of resources or interest in conducting active enforcement can affect the impact and duration of enforcement efforts.¹¹³

Judicial nullification of penalties directed at retailers was noted as a potential barrier in achieving and maintaining retailer compliance through active enforcement.^{89,114} Other reviews have suggested that replacing criminal offense statutes with specified civil penalties (e.g., graduated fines or license suspension) would improve enforcement efforts and minimize court appearances.^{2,84,114}

Another barrier to implementation can come from legislative efforts to weaken, replace, or prevent the implementation and conduct of interventions to reduce minors' access. Several reports identified in this review summarized these efforts.^{2,80,115,116}

In conclusion, the Task Force recommends community mobilization combined with additional interventions—such as stronger local laws directed at retailers, active enforcement of retailer sales laws, and retailer education with reinforcement—on the basis of sufficient evidence of effectiveness in reducing youth tobacco use and access to tobacco products from commercial sources. Preemption is a significant barrier to the implementation of the intervention combinations evaluated in this report. The published literature describes a number of legislative efforts to weaken, replace, or prevent the implementation and conduct of these interventions.

We also evaluated the effectiveness of several interventions implemented alone in reducing minors' access to and use of tobacco products. The evidence was insufficient to determine the effectiveness of any of these interventions by itself.

Sales Laws Directed at Tobacco Retailers to Reduce Illegal Sales to Minors When Implemented Alone: Insufficient Evidence to Determine Effectiveness

In addition to general laws governing the sale of tobacco, laws directed at retailers provide specific regulation or restriction of the sale of tobacco products to minors. These laws include licensing requirements for tobacco retailers and bans or restrictions on tobacco product vending machines and self-service displays. The laws may include additional conditions, such as requiring proof of the purchaser's age before selling tobacco, displaying sales laws (such as warning signs at the point of purchase), banning the sale of single cigarettes, and restricting the age of the seller.

These laws may designate the method of enforcement and establish the penalties and the responsible parties for each violation (for example, civil penalties directed at the retail owner or license holder). The interventions described in the studies we reviewed were implemented by local governments and were not limited by state preemption legislation. (The effectiveness of active enforcement of these laws is reviewed separately in this chapter.)

Effectiveness

- We found insufficient evidence to determine the effectiveness of sales laws directed at retailers, when implemented alone, in reducing minors' access to and use of tobacco products.
- Only one study qualified for review; it had limitations in study execution and lacked measurements of youth tobacco use or purchase behaviors.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review are based on a single qualifying study.¹¹⁷ Four additional reports^{24,32,118,119} (on three studies) were identified but did not meet our quality criteria and were excluded from the review. The single qualifying study evaluated the effect of a local county ban of self-service displays on sales of tobacco products to youth. No measurements of differences in tobacco use among youth were attempted. A slight reduction (3.2 percentage points from a baseline of 17.5%) in the number of retailers willing to sell tobacco to youth was observed. Results from this single qualifying study provided insufficient evidence to determine the effectiveness of sales laws directed at retailers when implemented alone in reducing youth access to tobacco.

Because we could not establish the effectiveness of these laws, we did not examine situations in which these laws would be applicable, information about economic efficiency, or possible barriers to implementation.

A potential benefit of these laws is that license requirements for the sale of tobacco products enable communities to identify commercial sources of tobacco. These laws also provide support for additional interventions, such as active enforcement of sales laws. Finally, self-service display bans reduce or eliminate minors' ability to obtain cigarettes by stealing them. No harms were identified.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of laws directed at retailers when implemented alone in reducing youth access to tobacco, because only one study qualified for our review, and that study had limitations in execution and lacked measurements of youth tobacco use or purchase behaviors.

Laws Directed at Minors' Purchase, Possession, or Use of Tobacco Products When Implemented Alone: Insufficient Evidence to Determine Effectiveness

These laws prohibit the purchase, possession, or use of tobacco products by minors. Communities have implemented laws directed at minors, and state governments are increasingly doing so. The laws may designate which agency is responsible for enforcement as well as the penalty for violations. Some laws require minors who have received citations to participate in educational programs that provide assistance in quitting smoking.

Effectiveness

- We found insufficient evidence to determine the effectiveness of laws directed at minors' purchase, possession, or use of tobacco products when implemented alone in reducing minors' access to and use of tobacco products.
- The evidence was insufficient because we identified no studies that met the quality criteria for our review.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

Our search identified no studies that evaluated the effectiveness of laws directed at minors' purchase, possession, or use of tobacco products when implemented alone. We therefore found insufficient evidence to determine whether or not these laws are effective in reducing minors' access to tobacco.

Because we could not establish the effectiveness of these laws, we did not examine situations in which they would be applicable, information about economic efficiency, or possible barriers to implementation.

Although we could not determine whether these laws are effective, we did note other published effects of the laws. Preemption legislation and policies in some states have directed resources and priority of enforcement away from laws directed at retailers in favor of laws directed at youth.^{114,120} Some laws establish or mandate participation in programs providing education and cessation assistance to cited youth.¹²¹

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of laws directed at minors' purchase, possession, or use of tobacco products when implemented alone, because no studies qualified for the review.

***Active Enforcement of Sales Laws Directed at Retailers When Implemented Alone:
Insufficient Evidence to Determine Effectiveness***

Active enforcement is used to achieve and maintain retailer compliance with sales laws. This enforcement consists of periodic unannounced compliance checks, which employ recruited, trained, and supervised minors to make standardized attempts to purchase tobacco products from retailers and vending machines. Retailers who violate sales laws receive citations from law enforcement officers or officers in designated government agencies (e.g., health department inspectors).

Local enforcement operations identified in this review were not limited by state preemption legislation.

Effectiveness

- We found insufficient evidence to determine the effectiveness of active enforcement of sales laws when implemented alone in reducing minors' access to and use of tobacco products.
- Evidence was insufficient because only one qualifying study was identified, which did not measure differences or changes in youth tobacco use or purchase behaviors.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review are based on one study.¹²² An additional study was identified but did not meet our quality criteria and was excluded from the review.¹²³

The single qualifying study evaluated the effect of an enforcement campaign on the sale of single cigarettes by tobacco retailers in New York City. Starting from a 100% baseline, the percentage of retailers willing to sell cigarettes in the intervention (enforcement) group decreased by 47.7 percentage points (singles or packs) on youth test purchases (the post-intervention sales

rate was 46.9%). This study did not measure differences or changes in youth tobacco use or purchase behaviors. We therefore found insufficient evidence to determine if active enforcement of existing laws against sale of tobacco products to minors, when implemented alone, is effective in reducing such sales.

Because we could not establish the effectiveness of these laws, we did not examine situations in which they would be applicable, information about economic efficiency, or possible barriers to implementation.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of active enforcement of sales laws when implemented alone in reducing youth access to tobacco, because only one qualifying study was identified and it did not provide measurements of youth tobacco use or purchase behaviors.

Our review identified two types of education interventions directed to tobacco retailers. In both interventions, the messages emphasized the importance of refusing to sell tobacco products to minors. These education interventions differed in the intensity and content of the educational messages, and in whether or not they provided reinforcement to retailers (in the form of follow-up and feedback).

Retailer Education with Reinforcement and Information on Health Consequences When Implemented Alone: Insufficient Evidence to Determine Effectiveness

These interventions aim to increase retailer compliance with prohibitions on tobacco sales to minors through repeated educational messages and feedback on retailer performance. Reviewed interventions involved face-to-face delivery of education messages by concerned citizens, health department workers, or law enforcement officers. The educational component included follow-up that provided either positive or negative reinforcement of retailer or clerk compliance with sales laws based on periodic unannounced compliance checks. In most cases, these interventions included information on the health consequences of tobacco use.

Effectiveness

- We found insufficient evidence to determine the effectiveness of retailer education providing reinforcement and information on health consequences when implemented alone in reducing minors' access to and use of tobacco products.
- The evidence was insufficient because we found no studies that met our quality criteria for review.

- Insufficient evidence means that we were not able to determine whether or not the intervention works.

Our search identified two studies in which retailer education with reinforcement was implemented alone.^{124,125} Both studies were excluded from the review, however, because one had limitations in the quality of execution and the other had a least suitable study design. Additionally, neither study evaluated differences or changes in youth tobacco use or purchase behaviors. Therefore, we found insufficient evidence to determine whether or not retailer education with reinforcement and information on health consequences, by itself, is effective in increasing retailer compliance with laws prohibiting the sale of tobacco products to minors.

Because we could not establish the effectiveness of this intervention, we did not examine situations in which such programs would be applicable, information about economic efficiency, or possible barriers to implementation.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of retailer education providing reinforcement and information on health consequences when implemented alone in reducing minors' access to tobacco, because no studies met our quality criteria for inclusion in the review.

***Retailer Education without Reinforcement When Implemented Alone:
Insufficient Evidence to Determine Effectiveness***

These education interventions deliver messages to retailers about tobacco sales to minors without providing follow-up and feedback on retailer performance. Education interventions, which aim to increase retailer compliance with prohibitions on the sale of tobacco to minors, include distribution of information about current or recent changes in local or state laws governing sales to minors as well as distribution of materials such as display signs or training manuals for clerks. In the studies evaluated in this review, education was carried out by mail; in face-to-face encounters; or through training sessions conducted by concerned citizens, health department workers, or law enforcement officers. The educational messages did not address the health consequences of tobacco use. Our review did not identify any studies evaluating mandatory training sessions for retailers or clerks or provisions to reduce or eliminate penalties for retailers who participated in education programs.

Effectiveness

- We found insufficient evidence to determine the effectiveness of retailer education without reinforcement when implemented alone in reducing minors' access to and use of tobacco products.

- Only three studies qualified for our review, and none of these measured youth tobacco use or purchase behaviors.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review are based on three studies.^{122,126,127} Three additional studies were identified but did not meet our quality criteria and were excluded from the review.^{128–130} The three reviewed studies evaluated mailed educational messages or a combination of mailed messages and face-to-face encounters. Although one study observed an increase (+21.3 percentage points) in the number of retailers requiring proof of age from youth making test purchase attempts and two studies observed absolute reductions in retailer sales rates of 9 and 27.5 percentage points, respectively, none of the studies provided measurements of differences or changes in youth tobacco use or purchase behaviors. Therefore, we were unable to determine the effectiveness of retailer education without reinforcement, by itself, in increasing retailer compliance with laws prohibiting the sale of tobacco products to minors.

Because we could not establish the effectiveness of the intervention, we did not examine situations in which it would be applicable, information about economic efficiency, or possible barriers to implementation.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of this intervention because of the small number of qualifying studies and lack of measurement of youth tobacco use or purchase behaviors.

Community Education about Minors' Access to Tobacco Products When Implemented Alone: Insufficient Evidence to Determine Effectiveness

These interventions attempt to disseminate information community-wide to focus public attention on the issue of youth access to tobacco products. The educational components include community-wide assessments of compliance by tobacco retailers, with results disseminated through mass media events, news coverage, and presentations to civic groups and local governments. The interventions can also include community and school meetings and activities as well as direct contact with local governments through testimony, petitions, letters, and phone calls.

In this review, we distinguished between community education efforts (reviewed here) and community mobilization, in which the education efforts fostered or were coordinated with additional interventions directed at minors' access to tobacco products (see Reducing Tobacco Use Initiation—Community Mobilization Combined with Additional Interventions).

Effectiveness

- We found insufficient evidence to determine the effectiveness of community education when implemented alone in reducing minors' access to and use of tobacco products.
- We identified no studies that evaluated this intervention.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

Our search identified no studies in which community education interventions were implemented and evaluated alone. We therefore had insufficient evidence to determine whether community education, by itself, is effective in increasing community awareness of the issue of youth access to tobacco products.

Because we could not establish the effectiveness of this intervention, we did not examine situations in which it would be applicable, information about economic efficiency, or possible barriers to implementation.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of this intervention in reducing minors' access to tobacco, because no qualifying studies of community education alone were identified.

Increasing Tobacco Use Cessation

Interventions to increase the number of tobacco users who successfully quit include efforts to increase the number of people who attempt to quit, efforts to improve the success rate for quit attempts, and efforts that support both of these goals. We reviewed two approaches appropriate for communities (increasing the unit price for tobacco products and mass media education); five approaches that can be implemented in healthcare systems (provider reminder systems alone; provider education programs alone; provider reminder systems and provider education programs together, with or without client education materials; provider feedback systems; and reducing client out-of-pocket costs for effective cessation therapies); and one intervention appropriate for both communities and healthcare systems (telephone cessation support).

In conducting this review of the evidence, the Task Force noted that spontaneous and unassisted rates of tobacco use cessation among tobacco users are low (3%–10%).^{6,131} Although the interventions reviewed in this section showed relatively small effects (increases of 2.2 to 4.1 percentage points), they represent relatively large improvements in the success rates for tobacco use cessation. Task Force assessments and conclusions of the evidence on effectiveness reflect these aspects of tobacco use cessation.

***Increasing the Unit Price for Tobacco Products:
Recommended (Strong Evidence of Effectiveness)***

Excise tax increases at the municipal, state, or federal level can raise the unit price for tobacco products. In several states, excise tax increases on tobacco products resulted from successful state ballot initiatives. Although other factors affect tobacco product pricing, excise tax increases have historically resulted in an equivalent or larger increase in tobacco product prices.²²

Effectiveness

- Increases in tobacco product price are consistently effective in reducing tobacco use (a 10% price increase results in approximately a 4% decrease).

Applicability

- These findings should be applicable to most adults in the United States.

Other Effects

- Price increases also reduce tobacco use among adolescents and young adults.

Barriers

- Increasing the excise tax on tobacco products requires passage of legislation or a statewide referendum.

The findings of our systematic review are based on 17 aggregate studies, which were consolidated from 51 papers that evaluated the effectiveness of price increases for tobacco products in reducing tobacco use.^{23,29,30,33,132-178} Because many of these papers analyzed the same data during similar, identical, or overlapping periods of time, we decided to consolidate papers into aggregate studies on the basis of similarities in location, the period of study, and the dataset employed. Five additional studies were identified but did not meet our quality criteria and were excluded from the review.¹⁷⁹⁻¹⁸³

The 17 aggregate studies reviewed evaluated the effect of tobacco product price on tobacco use in California; Massachusetts; Oregon; 11 Western states; national evaluations conducted in the 1990s, 1980s, and 1970s; and three studies conducted in Canada, three in the United Kingdom, and one each in Austria, Finland, Switzerland, and New Zealand.

Price elasticity of demand (the percentage change in consumption that results from a 1% change in price) was the most common measure used in these studies to estimate the effect of tobacco product price increases. Of the 17 aggregate studies, 13 included measurements of price elasticity of demand, which generally showed decreases in consumption. For example, in 10 aggregate studies that measured cigarette sales data, a 10% increase in product price would result in a 4.1% decrease in population consumption (range,

2.7% to 7.6%). Similar effects were seen in all measures of changes in tobacco prevalence and amount smoked.

Overall, this body of evidence documented consistent effectiveness of increases in tobacco product price in reducing tobacco use, regardless of the measurements reported or calculated, the setting or period of time evaluated, or differences in the control of potential confounders.

Four studies using measures other than price elasticity of demand (for example, measurements of tobacco product sales or consumption) showed that tobacco use declined in response to price increases, whereas excise tax decreases significantly slowed this decline.

These results should be applicable to most adults in the United States. All of the reviewed studies evaluated the effect of price on the consumption of cigarettes. Some also evaluated the effect of price increases on consumption of smokeless tobacco, cigars, and pipe tobacco. Studies demonstrated the effectiveness of state excise tax increases and federal excise tax increases in reducing tobacco consumption; effectiveness was also shown for whites, African Americans, Hispanics, men and women, across most social classes, and among people with incomes below median or with less than a high school education.

A positive effect of increasing the unit price for tobacco products is a decrease in tobacco use among adolescents and young adults (discussed under Reducing Tobacco Use Initiation). One potential negative effect of increases in tobacco product excise taxes is an increase in smuggling (illegal cross-border transport and sale of untaxed tobacco products). No recent studies of the effects of organized smuggling in the United States were identified in this review, however, and an analysis in 1985 reported a significant reduction in activity following federal legislation in 1978.¹⁸⁴

Effects of tobacco product price increases that might reduce (but would not eliminate) the potential health benefits of more cessation and less consumption include legal individual cross-border purchases of tobacco products; substitution of tobacco products (e.g., smokeless tobacco for cigarettes) created by unequal taxation on different kinds of tobacco products; and modification of individual tobacco use patterns, such as smoking cigarettes longer or changing to a higher-tar, higher-nicotine brand.¹⁴⁸

Econometric analyses were included in the studies qualifying for review of effectiveness. A separate economic evaluation was not conducted for this intervention review.

Passage of legislation or a statewide referendum is needed to increase the excise tax and may therefore present a barrier to implementation. Political op-

position has historically been well organized and funded at the municipal, state, and federal levels. Reports on both successful and unsuccessful state initiatives that proposed an increase on tobacco product excise taxes have been published.^{34–36}

In conclusion, the Task Force recommends increasing the unit price for tobacco products on the basis of strong evidence of effectiveness in increasing the number of people who stop using tobacco. Raising the unit price for tobacco products, through increases in excise taxes for tobacco products, is consistently effective in reducing tobacco use, regardless of the measure used. The findings of this review should be applicable to most adults in the United States. Tobacco product price increases also reduce tobacco use among adolescents and young adults. The need for passage by a state legislature or a referendum can be a significant barrier to efforts to raise excise taxes on tobacco products.

Mass Media Education

Mass media messages (broadcast and print) are disseminated to provide cessation information, motivation to assist tobacco users in their efforts to quit, or both. In our review, we distinguished among three subtypes of mass media interventions—campaigns, cessation series, and cessation contests—which differ in the duration, intent, and intensity of the media messages. Each is addressed separately below.

Mass Media Education Campaigns Combined with Other Interventions: Recommended (Strong Evidence of Effectiveness)

Mass media campaigns employ brief, recurring messages over time (weeks to years) to provide information or motivation to tobacco users and others (family members, households, peers) with the goal of increasing or improving efforts to quit using tobacco products. The messages, developed through formative research, are disseminated through paid broadcast time and print space, donated time and space (as public service announcements), or a combination of the two. Campaigns can focus on cessation or can include cessation themes within a broader range of tobacco messages (including messages directed at reducing tobacco use among youth, reducing secondhand smoke exposure, or both). Mass media campaigns can be combined with other interventions, such as an excise tax increase or additional community-wide education efforts.

Effectiveness

- Mass media education campaigns, when combined with other activities, were effective in increasing tobacco use cessation by approximately 2 additional quitters per 100.
- These interventions were also effective in reducing tobacco use in the population, as measured in states and in communities. Changes in tobacco use include reductions in consumption (by approximately 12.8%) and reductions in the prevalence of tobacco use (by approximately 3 people per 100 tobacco users).

Applicability

- These findings should be applicable to most people in the United States.

Other Effects

- Mass media education campaigns can increase the number of people who seek help from telephone “quit” lines.

The findings of our systematic review are based on 15 studies that evaluated the effectiveness of mass media campaigns in reducing tobacco use.^{138–140,185–196} Another 14 papers provided information on studies already included in the review.^{75,135,197–208} Three studies measured the effect of a mass media campaign in increasing use of a telephone cessation information service: these results are described below (see Multicomponent Interventions that Include Client Telephone Support).^{209–211} Six studies did not meet our quality criteria and were excluded from the review.^{212–217} All of the reviewed studies evaluated the combination of a mass media education campaign coordinated or concurrent with at least one other intervention: an excise tax increase (6 studies), community education programs such as the distribution of self-help cessation information (12 studies), individual or group cardiovascular disease risk factor reduction or smoking cessation counseling (7 studies), and other mass media efforts (2 studies).

The reviewed studies measured changes in tobacco use in terms of changes in individual tobacco use cessation, changes in population consumption (measured by statewide sales of cigarettes), and changes in the prevalence of tobacco use in the study population. In general, the studies observed increases in tobacco use cessation and decreases in population consumption and prevalence.

Five studies evaluated the effect of combined interventions including mass media campaigns on tobacco use cessation among groups of recruited tobacco users. In study periods of one to five years’ duration, the median difference in cessation between intervention group participants and comparison group participants (who were potentially exposed to the media component in

three studies) was an increase of 2.2 percentage points (representing approximately 2 additional people per 100 who quit using tobacco; range, 2 to 35).

Three studies, with follow-up periods of two to eight years, evaluated the effect of statewide mass media campaigns when combined with additional interventions (excise tax increase, community and school education activities) on consumption of cigarettes as determined by sales of taxed cigarette packs. In all three states, overall tobacco consumption declined at a greater rate than in the rest of the United States (median 12.8% decrease).

Finally, seven studies measured changes in tobacco use prevalence in communities exposed to a mass media campaign combined with additional interventions. Six studies observed decreases in tobacco use prevalence over study periods of 6 months to 20 years, and five of these studies included a concurrent comparison population. In these studies, tobacco use prevalence decreased in the intervention population by a median of 3.4 percentage points (range, -7 to +0.2) when compared with the unexposed community.

The results of these studies provide strong evidence of the effectiveness of mass media education campaigns, when combined with other activities, in increasing tobacco use cessation and reducing tobacco consumption.

These findings should be applicable to people throughout the United States. All of the reviewed studies focused on cigarette use. These studies included statewide and regional campaigns in the United States, national campaigns in Scotland, and regional campaigns in Finland. Studies were conducted in large cities as well as in smaller communities. In the United States, studies were also conducted specifically among Hispanics and among Vietnamese men.

Mass media campaigns, when combined with additional interventions, can also be effective in reducing tobacco use among youth (see Reducing Tobacco Use Initiation), especially when the campaign includes a variety of targeted messages and broadcast times. In California, Oregon, and Massachusetts, for example, campaigns included messages for children and adolescents as well as messages about the health effects of exposure to environmental tobacco smoke. Several studies noted significant increases in the use of telephone cessation information or support services ("quit" lines) when the media messages told audiences about these services.

The findings of our systematic review of economic evaluations of community-wide interventions to increase cessation among adult tobacco users are based on two cost-effectiveness studies (one conducted in the Netherlands²¹⁸ and the other in Scotland²¹⁹). Both studies reported program costs per quitter. In addition, the study in Scotland reported program cost per life-years saved, which was converted to dollars per QALY based on preference weights provided by Fiscella and Franks (a standard economic approach used to ad-

just health-related quality-of-life measures on a continuum from 0.0 [death] to 1.0 [optimal health]).²²⁰

The intervention conducted in the Netherlands evaluated the effect of a mass media campaign, a self-help manual, a hotline, and a nine-session group cessation program. Self-reported seven-day abstinence was determined at six-month follow-up. Costs included were wages, overhead, calls to the hotline, participant time, transportation, and charges for the group session. The effect size was estimated as the difference in smoking prevalence before and after the intervention. The self-reported cessation rate was 11% for participants using the self-help manual alone and 22% for participants using the self-help manual combined with the cessation program. Based on this effect size, adjusted program cost per quitter ranged from \$796 to \$1593.

The study conducted in Scotland consisted of a mass media campaign, telephone help line, and information booklet. At the 12-month follow-up, 9.8% of program participants reported having quit smoking for at least six months. Costs included research, production, design fees, printing, dissemination, staff salaries, and overhead. The adjusted program cost per quitter in this study ranged from \$298 to \$655. Adjusted program cost per QALY ranged from \$151 to \$328. The range was based on lower and upper bound estimates of adult participants.

The primary barrier to the implementation of mass media campaigns is the funding needed to develop and maintain an extended-duration, high-intensity campaign using paid and targeted broadcast times to deliver messages that resonate with target audiences. Cooperation between tobacco control programs can reduce program development costs. The barriers encountered by the tobacco control program in the state of California provide important lessons on the need for both public and political support and vigilance in maintaining an effective campaign.^{221,222}

In conclusion, the Task Force recommends mass media education campaigns combined with additional interventions on the basis of strong evidence of effectiveness in increasing tobacco use cessation and in reducing tobacco consumption. The findings of this review should be applicable to most people in the United States. An additional benefit of these campaigns can be an increase in the number of people who call in to dedicated telephone cessation support services' quit lines.

Mass Media Education—Cessation Series: Insufficient Evidence to Determine Effectiveness

Cessation series consist of broadcasted instructional segments designed to recruit, inform, and motivate users of tobacco products to try quitting and to succeed. Cessation series can be coordinated with pre-series broadcast or

print promotion, community education (e.g., distribution of self-help cessation materials), and organization of cessation groups in the community. The series can extend for a period of several weeks to several months. Techniques include nightly or weekly segments on news or informational programs giving expert advice or sharing peer group experiences on cessation issues (e.g., dealing with the symptoms of withdrawal). The series can encourage tobacco users to join a community-wide quit effort. Over the course of the broadcasts, viewers can receive ongoing support and assistance from cessation experts and recruited peers.

Effectiveness

- We found insufficient evidence to determine the effectiveness of mass media education cessation series in increasing the number of people who quit using tobacco products.
- Evidence was insufficient because of inconsistent results and inadequate comparison groups.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review are based on nine studies that evaluated the effectiveness of cessation series in increasing tobacco use cessation.^{223–231} Eleven additional studies were identified but did not meet our quality criteria and were excluded from the review.^{208,232–241} Ten other reports provided information on studies already included in the review.^{242–251}

All of the reviewed studies evaluated the effectiveness of cessation series combined with other interventions, such as community education (typically, access to or distribution of self-help cessation manuals), organized cessation groups or programs, or telephone cessation support (quit lines). Eight studies evaluated televised cessation series broadcast over periods of 20 days to three months, and one study evaluated a week-long newspaper cessation series. Five of the nine studies in the review evaluated broadcast cessation series conducted in the Chicago metropolitan area (in one of three waves) between 1985 and 1987. These studies were evaluated separately because of differences in the study populations or settings.

The reviewed studies provided insufficient evidence to determine the effectiveness of mass media education cessation series in increasing tobacco use cessation. The differences in cessation observed in these studies might be the result of (1) baseline differences in motivation to quit between intervention and comparison smokers and (2) the small group cessation sessions provided to the intervention group participants.

Because we could not establish the effectiveness of mass media cessation series in increasing the number of people who successfully quit using tobacco

products, we did not examine situations in which they would be applicable, information about economic efficiency, or possible barriers to implementation.

No harms or benefits of mass media cessation series were identified.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of mass media cessation series in increasing the number of people who successfully quit using tobacco products because of inconsistent results and inadequate comparison groups.

Mass Media Education—Cessation Contests: Insufficient Evidence to Determine Effectiveness

Cessation contests are community-wide events of short duration that use mass media to recruit and motivate users of tobacco products to participate in a program to quit by a certain date or during a specified time period. Cessation contests use both mass media and such small media as posters and flyers as the primary tool for promotion and to recruit tobacco product users in the community. Contests can increase cessation in the community by changing tobacco product users' attitudes about cessation, recruiting users to initiate a quit attempt, and motivating those who attempt to quit to remain abstinent through incentives or by mobilizing support from family, friends, and other participants. We evaluated contests that offered additional incentives for participation and successful cessation, as well as targeted quit events conducted without additional incentives.

Effectiveness

- We found insufficient evidence to determine the effectiveness of mass media education cessation contests in increasing the number of people who quit using tobacco products.
- Evidence was insufficient because only one study met the quality criteria for this review, and that study lacked sufficient evidence on which to base a recommendation.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review are based on one study.²⁵² We identified another 16 studies that did not meet our quality criteria and were excluded from the review.^{234,253–267} Three other papers provided information on the study already included in the review.^{268–270}

The single reviewed study evaluated a multicomponent smoking cessation program in New York City. Interventions included a cessation manual and video, telephone cessation support, and the opportunity to participate in smoking cessation contests.

Although the single reviewed study showed some improvements in self-reported cessation at a six-month follow-up, the evidence was insufficient to determine the effectiveness of mass media cessation series in increasing the number of people who successfully quit using tobacco products because the evidence in this single study was not sufficient to support a recommendation.

Because we could not establish the effectiveness of mass media cessation contests in increasing the number of people who successfully quit using tobacco products, we did not examine situations in which the programs would be applicable, information about economic efficiency, or possible barriers to implementation.

No harms or benefits of mass media cessation contests were identified.

In conclusion, evidence was insufficient to determine the effectiveness of mass media education cessation contests in increasing the number of people who successfully quit using tobacco products because only a single study, showing only moderate improvement, qualified for review.

Provider- and System-Based Interventions

We reviewed six interventions that can be implemented by healthcare systems and providers to increase cessation of tobacco use by clients. We did not evaluate the effectiveness of provider counseling to tobacco-using clients or the effectiveness of specific clinical therapies, which have been thoroughly reviewed by others.^{2,6,7,17}

Healthcare Provider Reminder Systems: Recommended (Sufficient Evidence of Effectiveness)

Provider reminder systems can identify clients who use tobacco products and prompt providers to discuss cessation with their clients or advise clients to quit at every encounter. Because even brief provider advice has a demonstrated effect on getting clients to quit,^{6,17} increasing the delivery of advice by providers should increase the number of clients who quit.

Reminders can be delivered by a variety of methods, including chart stickers, vital sign stamps, medical record flowsheets, and checklists, and the content of the reminders can vary. Provider reminder systems are often combined with other interventions, such as provider education and client education. Here we review provider reminder systems used alone; multicomponent interventions including provider reminders are considered below (see Healthcare Provider Reminder Systems with Provider Education, with or without Client Education).

Effectiveness

- Provider reminder systems were effective in increasing the number of clients who quit smoking by approximately 4 additional clients per 100.
- Provider reminder systems were also effective in increasing the number of clients whom providers advise to quit smoking by approximately 13 additional clients per 100.
- These systems were also effective in increasing the determination of client smoking status by providers by approximately 32 additional clients per 100.

Applicability

- These findings should be applicable to most clinical setting in the United States.

Other Effects

- The use of provider reminder systems may also increase delivery of other preventive services.

The findings of our systematic review are based on seven studies evaluating the effectiveness of provider reminder systems when used alone.²⁷¹⁻²⁷⁷ One additional paper was identified but did not meet our quality criteria and was excluded from the review.²⁷⁸ The reviewed studies prompted providers with chart prompts or stickers, “expanded vital signs” that indicate whether or not a client uses tobacco, and flowsheets. In one study, analysis was based on receipt of a consultation that encouraged implementation of a provider reminder system, not on actual implementation of the reminder system.

The seven reviewed studies used diverse measures to document changes in client smoking status, provider delivery of advice to quit, and client smoking cessation. The median improvement in determining client smoking status was 32.5 percentage points (range, 26 to 57.6; four studies); that is, the intervention resulted in determining the smoking status of approximately 32 additional clients per 100, with results ranging from 26 to 58 additional clients per 100. These results were measured 8 to 24 months after implementation (median, 15 months). For provider delivery of advice to quit, a median change of 13 percentage points was observed (range, 7 to 31; five studies) in assessments extending 2 to 24 months after implementation (median 8 months). One study reported a difference in biochemically confirmed smoking cessation of 4 percentage points six months after a clinic visit. These findings provide sufficient evidence that healthcare provider reminder systems are effective in increasing the number of tobacco-using clients who receive advice to quit from their providers.

These findings should be applicable in most clinical settings in the United States. Studies were conducted in primary care clinics, family practice clin-

ics, and pulmonary clinics. Most studies did not provide demographic information about client populations.

Two studies^{274,276} evaluated provider reminder systems that included preventive services in addition to delivering advice to quit to tobacco-using clients. Improvements were observed in some or all of the prompted activities. No other positive or negative effects were identified in this review.

We did not find any economic evaluations of provider reminder systems.

Although one potential barrier to the implementation of a provider reminder system could be the administrative burden, this was not identified as a problem in any of the reviewed studies, and most of the reminder systems (e.g., “expanded vital signs”) were easily implemented.

In conclusion, the Task Force recommends provider reminder systems on the basis of sufficient evidence of effectiveness in increasing the number of tobacco-using clients whom providers advise to quit (because this outcome has been previously shown to increase subsequent tobacco use cessation¹⁷). These findings should be applicable to most clinical settings in the United States. An additional benefit of provider reminder systems can be increased delivery of other preventive services.

Healthcare Provider Reminder Systems with Provider Education, with or without Client Education: Recommended (Strong Evidence of Effectiveness)

These multicomponent interventions to increase tobacco use cessation include efforts to educate and to prompt providers to identify and to intervene with tobacco-using clients, as well as to provide supplementary educational materials when appropriate. The interventions consist of a provider reminder system and a provider education program, and may or may not include client education materials (e.g., self-help cessation manuals).

A multicomponent intervention can provide an integrated approach to increasing and improving tobacco use cessation by clients. Goals of the interventions include educating, motivating, and prompting providers to increase and improve their interaction with tobacco-using clients, as well as improving client cessation by increasing knowledge and motivation to quit and to remain abstinent. The multicomponent interventions evaluated in this section include at least one provider-directed component.

Effectiveness

- These interventions were effective in increasing the number of clients who quit smoking by approximately 5 additional clients per 100.
- The interventions were also effective in increasing the number of tobacco-

using clients who received advice to quit from their healthcare provider by approximately 20 additional clients per 100.

Applicability

- These findings should be applicable to most clinical settings in the United States and to a variety of provider specialties.

Other Effects

- The use of provider reminder systems plus provider education, with or without client education, may also increase delivery of other preventive services.

The findings of our systematic review are based on 31 studies that evaluated the effectiveness of these interventions in reducing tobacco use among clients.^{272,275,279–307} One study measured changes in adolescent tobacco use initiation and is not considered further in this section.²⁹⁰ Fifteen additional papers provided information on studies already included in the review.^{308–322} Six additional studies were identified but did not meet our quality criteria and were excluded from the review.^{323–328}

Overall, studies that measured providers' delivery of advice to quit using tobacco and the number of clients who quit showed significant improvements. For example, the 20 studies that evaluated the effectiveness of a multi-component intervention combining at least a provider reminder system and a provider education program found a median change of 20 percentage points (range, 5 to 60) in the number of clients receiving advice to quit from their providers and a median change of 5 percentage points (range, -1 to +26) in the number of clients who quit using tobacco (in follow-up periods of five weeks to 12 months). Similar results were found in the subsets of studies that evaluated interventions using only a provider reminder system and a provider education program (seven studies); interventions that also included client education materials (13 studies); and interventions that included provider reminders and client education but not provider education (three studies). Interventions that included only provider education are discussed below (see Healthcare Provider Education Alone).

These results show that healthcare provider reminder systems plus provider education programs, whether or not combined with client education, are effective in increasing the number of tobacco-using clients who receive advice to quit from their healthcare providers.

These findings should be applicable to a variety of healthcare settings and provider specialties. Studies were conducted in health maintenance organizations (HMOs), private practices, academic health care centers, physician training programs, and public health clinics, among providers in primary

care, internal medicine, family medicine, obstetrics, pediatrics, and dentistry. One study focused on reducing use of smokeless tobacco.

Provider reminder systems can include prompts for additional preventive services. No other positive or negative effects were identified in the review.

We did not find any economic evaluations of these interventions.

The burden of administering provider reminder systems is a potential barrier to their implementation.

In conclusion, the Task Force recommends multicomponent interventions including provider reminder systems and provider education, with or without client education, on the basis of strong evidence of effectiveness in increasing the number of providers who advise their clients to quit smoking and in increasing the number of clients who quit. These findings should be applicable to most clinical settings in the United States and to relevant clinical specialties. Reminder systems can include prompts for additional preventive services.

Healthcare Provider Education Alone: Insufficient Evidence to Determine Effectiveness

The goal of provider education is to increase providers' knowledge about tobacco use and cessation, to change their attitudes and practices, and increase or improve their interactions with clients who use tobacco. These interactions can include identifying more tobacco-using clients, increasing delivery of advice to quit, improving the quality of advice to quit, and both increasing and improving providers' efforts to assist clients in their attempts to quit and to remain tobacco-free.

Information can be delivered in a variety of ways, including lectures, written materials, videos, and continuing medical education seminars. Physicians, nurses, physician assistants, students, and office staff can receive this education.

Provider education efforts are frequently combined with other interventions, such as provider reminders and client education efforts. Multicomponent interventions that include provider reminders are considered above (see Healthcare Provider Reminder Systems with Provider Education, with or without Client Education).

Effectiveness

- We found insufficient evidence to determine the effectiveness of provider education alone in increasing the number of people who quit using tobacco products.
- Evidence was insufficient because (1) few studies measured the effect of the intervention on tobacco use cessation and (2) the results of studies that

measured changes in the number of clients whom providers advise to quit smoking were inconsistent.

- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review are based on 16 studies that evaluated the effectiveness of provider education interventions when implemented alone.^{272,275,285,294,301,302,329–338} Two additional papers provided information on studies already included in the review.^{339,340} Nine other studies were identified but did not meet our quality criteria and were excluded from the review.^{327,341–348}

Provider education techniques evaluated in the reviewed studies included day-long seminars, lectures for practitioners and for resident physicians, lectures and office visits or contacts, small group tutorial sessions, mock interviews with feedback, and education materials. The provider education sessions ranged from two hours to three days (median, two and a half hours) in the 11 studies that provided this information.

The 16 qualifying studies reported a total of 19 measures of change in provider counseling skills or behaviors and 2 measures of client tobacco use. Of the 10 studies that measured differences in provider delivery of advice to quit, 4 observed either no effect or a negative effect (median change 2.2 percentage points; range, –5 to 73). Only two studies reported differences in client tobacco use cessation, showing small increases (1.7 and 5.2 percentage points, respectively). Five studies measured differences in provider determination of client smoking status, with percentage differences ranging from 0.1 to 35 percentage points (median, 8).

These results provide insufficient evidence to determine whether health-care provider education alone is effective in changing providers' knowledge, attitudes, and practices about tobacco use or in increasing the number of people who stop using tobacco products.

Because we could not establish the effectiveness of these interventions, we did not examine situations in which they would be applicable, information about economic efficiency, or possible barriers to implementation.

No harms or benefits of provider education alone were identified.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of provider education interventions when implemented alone in increasing the number of people who quit using tobacco products. Evidence was insufficient because (1) few studies evaluated the effect on client tobacco use cessation and (2) studies that evaluated provider delivery of advice to quit had inconsistent results.

Healthcare Provider Feedback and Assessment: Insufficient Evidence to Determine Effectiveness

Feedback interventions inform and motivate providers by assessing their performance in identifying client tobacco use status, delivering advice to quit, or both. Assessment techniques include chart reviews and the use of computerized records. These interventions can be combined with other activities, such as provider reminders and provider education, and these combinations were considered in our review.

Provider assessment and feedback can motivate providers to increase and improve their interactions with clients in such areas as advising clients to stop using tobacco. Evaluation of provider assessment and feedback is timely because (1) clinical information systems are improving and are increasingly common; (2) effective cessation therapies are available, and an increase in provider interactions with tobacco-using clients could increase the use of these therapies; and (3) quality assurance approaches such as the Health-plan Employer Data and Information Set (HEDIS) are being used more often.

Effectiveness

- We found insufficient evidence to determine the effectiveness of provider assessment and feedback in increasing either provider advice to clients to quit using tobacco or the number of clients who quit using tobacco products.
- Evidence was insufficient because the small number of available studies did not measure these outcomes.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review of the effectiveness of provider assessment and feedback interventions in changing provider behaviors toward tobacco-using clients are based on three studies.^{349–351} Two additional studies were identified but did not meet our quality criteria and were excluded from the review.^{352,353}

A provider assessment and feedback program was evaluated alone in one study, and in combination with other components in two studies that included a provider education program and a provider reminder flowsheet. In all three studies, provider documentation or recognition of a client's tobacco use status was only 1 of up to 26 preventive care practices for which assessment and feedback were provided. Only one study used a computer system to collect information and enable providers to obtain feedback information.

None of the qualifying studies attempted to measure changes in provider delivery of advice to quit or whether clients quit using tobacco. The three studies did, however, provide measures of effectiveness in increasing provider recognition of client tobacco use status, showing a median increase of

21 percentage points (range, 13 to 39) in study periods that ranged from three months to six years. This improvement, however, provides insufficient evidence to determine the effectiveness of healthcare provider feedback alone in changing providers' knowledge, attitudes, and practices about tobacco use or in increasing the number of people who stop using tobacco products.

Because we could not establish the effectiveness of these interventions, we did not examine situations in which the programs would be applicable, information about economic efficiency, or possible barriers to implementation.

An additional benefit of increased provider delivery of other preventive care practices was found in two studies. No other positive or negative effects were identified in this review.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of provider assessment and feedback interventions, when implemented alone, in increasing either provider advice to clients to quit using tobacco or the number of clients who quit using tobacco products. Evidence was insufficient because the small number of available studies did not provide measurements of provider advice to clients to quit using tobacco or the number of clients who quit using tobacco products.

***Reducing Client Out-of-Pocket Costs for Effective Cessation Therapies:
Recommended (Sufficient Evidence of Effectiveness)***

Reducing the financial barriers that may prevent clients from using effective cessation therapies such as nicotine replacement,⁶ other pharmacologic therapy,⁶ behavioral therapies such as cessation groups,⁶ or a combination of these approaches is intended to increase the use of effective therapies, increase the number of people who try to quit, and increase the number of people who succeed in quitting.

Effectiveness

- Reducing client out-of-pocket costs is effective in increasing the number of people who successfully quit using tobacco by approximately 8 clients per 100.
- The intervention is also effective in increasing the number of people who use cessation therapies by approximately 7 clients per 100.

Applicability

- These findings should be applicable in a variety of clinical settings in the United States, in rural and mixed rural-urban settings.

Other Effects

- Clients may use nicotine gum beyond the recommended length of time.

Barriers

- Insurance requirements to combine pharmaceutical therapies with behavioral programs may discourage some clients from trying to quit.

The findings of our systematic review are based on five studies that evaluated the effectiveness of reducing clients' out-of-pocket costs for nicotine gum or nicotine replacements.^{354–358} In two studies, nicotine gum or a nicotine replacement were provided as part of, or in addition to, a behavioral program. Access to a behavioral program was provided but rarely used in a third study. In three studies, nicotine gum or nicotine replacement was provided free of charge to participants in the intervention group. In one study, the out-of-pocket costs of the combination of behavioral program and nicotine replacement were \$10 for intervention group clients and \$52.50 for comparison group clients.

Overall, studies showed improvement in both the use of cessation therapies and the number of clients who quit using tobacco. For example, three studies reported a median difference in use of cessation therapies of 7 percentage points (range, 6.5 to 28). Four studies that measured changes in the number of people who stopped using tobacco showed a median increase of 7.8 percentage points (range, 2.1 to 11) in follow-up periods that ranged from 6 to 12 months (median, 9 months). These findings show that reducing client out-of-pocket costs for tobacco cessation therapies is effective in increasing both the use of these therapies and the number of tobacco users who quit.

These findings should be applicable to different settings and populations. Studies were conducted in the United States; in HMOs, private practices, and a Department of Defense hospital; in rural and mixed rural-urban settings; and among a low-income population.

A potential harm of nicotine replacement therapy was found in the extended use of nicotine gum beyond the recommended duration (four months) in one study. No other potential benefits or harms were identified in this review

The findings of our systematic review of economic evaluations are based on two studies.^{355,356} One study,³⁵⁵ a cost-effectiveness analysis that reported program cost per quitter, was conducted in a healthcare setting in Washington State for employees enrolled in a health plan. The intervention consisted of insurance coverage for clients in a behavioral program that included nicotine replacement, with a 12-month follow-up. Four types of insurance coverage, with different out-of-pocket costs to users, were available: (1) standard coverage: a 50% co-payment for the behavioral program and the usual \$5 co-

payment per prescription for nicotine replacement therapy; (2) reduced coverage: a 50% co-payment for both the behavioral program and nicotine-replacement therapy; (3) flipped coverage: no co-payment for the behavioral program but a 50% co-payment for nicotine-replacement therapy, and; (4) full coverage: no co-payment for the behavioral program and the usual \$5 co-payment per prescription for nicotine replacement therapy. Costs measured included drugs, personnel, and cost of the behavioral program. Development, marketing, and implementation of the coverage plan were not included in the analysis. The adjusted program costs per quitter were \$135, \$141, \$149, and \$195 for standard, reduced, flipped, and full coverage, respectively.

The second study³⁵⁶ was a cost-benefit analysis conducted in Vermont, reporting net benefit. The intervention, conducted at a rural family practice clinic with low-income clients, consisted of brief physician advice and a prescription for free nicotine gum, with a six-month follow-up. Costs measured included physician time, nicotine gum, smoking cessation booklets, and client time. Development, promotion, and evaluation costs were not included. The adjusted quit rate for the intervention group was 9.4%. When costs and benefits from averted illness were compared, the intervention was shown to be cost-saving.

Potential barriers to increased use of effective cessation therapies are created by coverage requirements that tie pharmacotherapy to behavioral therapy interventions. Recent reviews have identified each strategy as effective alone.⁶ The combinations demonstrate higher cost-effectiveness³⁵⁹ but also limit the use of effective therapies for smokers who are unwilling to participate in the behavioral program. Including proactive telephone counseling as a behavioral therapy option might reduce these barriers.³⁶⁰

In conclusion, the Task Force recommends reducing out-of-pocket costs on the basis of sufficient evidence of effectiveness in increasing the number of people who use cessation therapies and in increasing the number of people who successfully quit using tobacco. This approach has been tried successfully in a variety of clinical settings in both rural and rural-urban settings. A potential harm is that clients may continue to use nicotine gum beyond the recommended length of time. Healthcare coverage requirements for behavioral programs along with pharmaceutical therapies may discourage clients who do not wish to be part of the behavioral programs.

***Multicomponent Interventions That Include Client Telephone Support:
Recommended (Strong Evidence of Effectiveness)***

Telephone support for people trying to stop using tobacco provides counseling or assistance in quitting and in staying tobacco-free. Depending on the

program, tobacco users or healthcare providers can initiate contact. Telephone support can include trained counselors, healthcare providers, or taped messages in single or multiple sessions that usually follow a standardized protocol for providing advice and counseling. Telephone support is usually combined with other interventions, such as client education materials, individual or group cessation counseling, or nicotine replacement therapies.

Telephone contact is intended to increase the motivation of tobacco users to attempt to quit, and can help to reduce relapses by providing support and assistance to recent quitters. In community settings, telephone support typically provides access to self-help cessation materials and available local resources (such as group sessions), and may provide counseling and motivation sessions. In clinical settings, telephone follow-up calls usually support other clinical cessation interventions such as provider counseling, group cessation sessions, or nicotine replacement or other therapies.

Effectiveness

- These multicomponent interventions were effective in increasing the number of people who stop using tobacco by approximately 3 people per 100.

Applicability

- These findings should be applicable to most adults in a variety of clinical and community-wide settings in the United States.

The findings of our systematic review are based on 32 studies that evaluated the effectiveness of telephone cessation support.^{248,285,288,289,293,297,300,304,361-384} Sixteen additional papers provided information on studies already included in the review.^{309,310,312,315-319,321,385-391} Seven additional studies were identified but did not meet our quality criteria and were excluded from the review.^{216,328,392-396}

In all of the reviewed studies, telephone support was coordinated with additional interventions including client education (29 studies), provider-delivered counseling (17 studies), nicotine replacement (4 studies), a smoking cessation clinic (1 study), and a televised cessation series (1 study). Telephone support was reactive (the caller initiated all contact) in 5 studies and proactive (the provider initiated contact or the caller initiated contact, with provider follow-up) in 27 studies.

Thirty studies compared differences in tobacco use cessation based on use of or exposure to telephone support. In follow-up periods of 5 weeks to 34 months (median, 12 months) the median difference was an increase in tobacco use cessation of 2.6 percentage points (range, -3.4 to +23). Studies that compared telephone support plus client education to client education alone found similar increases in cessation. These findings show that multicomponent interventions that include client telephone support are effective in increasing tobacco use cessation.

These findings should be applicable in many settings and to many populations throughout the United States. Studies were conducted in HMOs, private practices, public health clinics, medical centers and hospitals, and resident training programs. Provider specialties included dentistry, obstetrics, gynecology and family planning, primary care, family practice, and internal medicine. Clients included hospitalized smokers, veterans, pregnant women, African Americans, and smokers over 60 years of age. One study focused on reducing the use of smokeless tobacco.

We found no other positive or negative effects of telephone cessation support.

The findings of our systematic review of economic evaluations are based on five cost-effectiveness studies.^{389,397-400} Two studies reported program costs per quitter,^{398,400} and three studies^{389,397,399} reported program costs per life-year saved ratios, which were converted to cost per QALY using preference weights reported by Fiscella and Franks (a standard economic approach used to adjust health-related quality-of-life measures on a continuum from 0.0 [death] to 1.0 [optimal health]).²²⁰

Both studies reporting program costs per quitter evaluated interventions to increase tobacco use cessation among pregnant women. The first study was conducted in Southern California in an HMO setting.³⁹⁸ The intervention consisted of a combined prenatal nutrition counseling and smoking cessation program aimed at reducing the incidence of low-birthweight infants. The smoking cessation program consisted of an eight-week home correspondence program that included weekly telephone calls to an automated answering service. Smoking cessation increased by 12 percentage points over the comparison group cessation rate of 38%. Costs included salaries, overhead, supplies, printing, phone, and postage. Adjusted program cost per quitter was \$677. The second study was a nationwide modeled intervention in the United States.⁴⁰⁰ It consisted of a single 15-minute counseling session conducted by a nonmedical counselor, instructional material, and two follow-up telephone calls. The change in cessation was an increase of 15 percentage points modeled from earlier randomized trials of smoking cessation among pregnant women. Costs included instructional materials, staff time, overhead, and training. The adjusted program cost per quitter was \$292. The difference in program cost between the two studies can be explained by the fact that the first study looked at a comprehensive intervention using more resources.

Of the three studies reporting program costs in terms of life-years saved, two studies looked at interventions conducted in hospital settings with adult clients who smoked. One study was conducted at the Mayo Clinic in Minnesota,³⁹⁷ and the other was conducted at HMO hospitals in Oregon and Washington.³⁸⁹ The comparison group quit rates for the studies were 10.7% and 9.2 %, respectively; the intervention group showed cessation increases

of 12 percentage points and 4 percentage points above the comparison quit rates, respectively. The Mayo Clinic intervention consisted of two programs: an individual nicotine-dependency treatment program and a relapse prevention program. The relapse prevention program included telephone follow-up calls, letters, and a mailed survey. The comparison group consisted of clients who received no program. Costs included personnel, supplies, telephone, drugs, and capital equipment. The adjusted program cost per QALY was \$2532. The HMO hospital intervention consisted of a 20-minute bedside counseling session, a video, self-help materials, and follow-up calls. Costs included program development, personnel, communications, and overhead. The adjusted program cost per QALY of this intervention was \$1248. The Mayo Clinic intervention had a higher program cost per QALY in spite of showing a higher net effect size, because programs costs of this intervention included letters, surveys, and treatment for nicotine dependency in addition to counseling and telephone calls. The third study,³⁹⁹ conducted at a Boston hospital with clients who had had an acute myocardial infarction, modeled an intervention consisting of nurse-managed smoking-cessation counseling including telephone support after discharge. The comparison group—clients exposed to standard smoking cessation counseling designed for survivors of acute myocardial infarction—had a quit rate of 45%. Cessation increased by 26 percentage points in the intervention group. Costs included were personnel and instructional materials. Time spent on the phone, follow-up time, program development, and training costs were not included. The adjusted program cost per QALY was \$73.

Lack of client awareness of, or interest in, support lines can limit the effectiveness of this intervention. Making clients aware of this available support (e.g., through media messages) is one way to increase use of support (quit) lines.

In conclusion, the Task Force Recommends telephone cessation support on the basis of strong evidence of effectiveness in increasing tobacco use cessation when implemented with other interventions (e.g., other educational approaches or clinical therapies) in both clinical and community settings. Effective interventions combined at least proactive telephone support and client cessation materials.

Reducing Exposure to Environmental Tobacco Smoke

Interventions to reduce exposure to environmental tobacco smoke (ETS) require or encourage the establishment of smoke-free areas in workplaces, in public areas, and in the home in an effort to reduce tobacco-related morbid-

ity and mortality. Having smoke-free areas can reduce exposure to ETS and ETS-related illness and death.^{12,13,401} Smoke-free policies can change the attitudes and behaviors of smokers and increase both the number of people who try to quit and the number of attempts made by each person. Smoke-free policies may also improve the success rate for each quit attempt by reducing opportunities for relapse.⁴⁰² Smoke-free policies also challenge the perception of smoking as a normal adult behavior.⁴⁰³ By changing this perception, these policies can change the attitudes and behaviors of adolescents, resulting in a reduction in tobacco use initiation.⁴⁰⁴

The interventions reviewed here are smoking bans and restrictions to address exposure in the workplace and in public areas and community education to reduce exposure to ETS, especially among children, in the home environment.

Smoking Bans and Restrictions: Recommended (Strong Evidence of Effectiveness)

Smoking bans and restrictions are policies, regulations, and laws established by private, nongovernment, and government groups and agencies. Smoking bans entirely prohibit smoking in geographically defined areas, whereas smoking restrictions limit smoking to designated areas. Smoking bans and restrictions can be implemented with additional interventions, such as education and tobacco use treatment programs.

Businesses establish smoking policies to protect employees and customers from exposure to ETS in the workplace. Accrediting agencies (e.g., the Joint Commission on Accreditation of Healthcare Organizations) set regulations to protect employees and patrons within their organizations. Federal, state, or local laws are implemented to protect people from ETS exposure in public areas, and to establish minimum standards for both public and private workplaces. Standards for regulations and laws establishing smoking restrictions often include the size, location, and ventilation requirements for designated smoking areas.

Effectiveness

- Smoking bans and restrictions were effective in decreasing the amount of ETS by approximately 72%.
- Bans and restrictions were also effective in reducing exposure to ETS by approximately 60%.

Applicability

- These findings should be applicable to most indoor workers in the United States.

Other Effects

- Smoking bans and restrictions also helped to reduce cigarette consumption and to increase the number of people who quit smoking.
- We found no adverse economic impacts on business or tourism as a result of these policies.

Barriers

- Preemption, the adoption or existence of a law at the state level that supercedes or precludes stronger local laws, can prevent implementation of smoking bans or restrictions.

The findings of our systematic review are based on 10 studies that evaluated the effect of smoking bans and restrictions on exposure to ETS.^{405–414} Seven additional studies were identified but did not meet our quality criteria and were excluded from the review.^{415–421}

The 10 studies provided a total of 12 measures of the effect of smoking bans and restrictions on exposure to ETS. Overall, 9 of the 10 studies observed reductions or differences in ETS exposure in workplaces that had smoking bans or restrictions. In four studies that measured environmental components of ETS (such as nicotine vapor) before and after implementation of the smoking ban or restriction, the median change, assessed 6 to 12 months after implementation of the ban or restriction, was a decrease of 72% (range, 44% to 97%). Six studies provided a total of eight measures of differences in self-reported exposure to ETS. In assessments conducted 4 to 18 months after implementation, the median change in self-reported ETS exposure was a decrease of 60% (range, 4% increase to 94% decrease).

Four studies evaluated the effect of smoking restrictions, four studies measured the effect of smoking bans, and two studies measured differences in workplace ETS exposure for both. In general, reductions in ETS exposure were greater in workplaces that had smoking bans than in those with only smoking restrictions.

These results show that smoking bans and restrictions are effective in decreasing both the amount of and exposure to ETS.

These results should be applicable to most indoor workers in the United States. Smoking bans and restrictions were evaluated in a variety of settings including hospitals and medical centers, healthcare provider offices, government or public sector workplaces, and a university. Studied bans and restrictions were created through a government law, private-sector policies, workplace policies, and local ordinance. Studies on representative samples of indoor workers in the states of California and Missouri, and on large, diverse samples of government employees in Texas and HMO employees in Oregon showed that smoking bans and restrictions reduced self-reported ETS expo-

sure in workplaces community-wide. No studies evaluated the effect of smoking bans or restrictions in public settings outside of the workplace, such as public transportation systems or sports and entertainment venues.

A number of additional benefits of smoking bans and restrictions were described in the studies in this review. Workers exposed to a workplace smoking ban or restriction reported a greater reduction in daily consumption of cigarettes (median, -1.2 cigarettes per day, nine qualifying studies^{409,410,412,422-427}) than did workers not exposed to a workplace smoking policy. Four studies of workplace smoking bans^{423,425-427} found that tobacco users exposed to the ban quit at a greater rate than did tobacco users who were not exposed to a ban in the workplace. Six studies⁴²⁸⁻⁴³⁴ (in seven reports) identified in this review evaluated the economic impact of smoking ordinances and found no adverse economic effects on businesses (including bars and restaurants) or on tourism.

The findings of our systematic review of economic evaluations are based on one study conducted in the United States, which modeled the costs and benefits of a proposed national smoke-free environment act to restrict or ban smoking inside all nonresidential buildings regularly entered by 10 or more people per week.⁴³⁵ Costs included implementation of the restriction or ban by the establishment, construction and maintenance of smoking lounges, and enforcement. Benefits included savings on medical costs by averting heart disease, the value of lives saved, costs averted by reduced smoking-related fires, and productivity improvements. The net present benefit to society (benefits minus costs) was in the range of \$42 to \$78 billion. This range was based on high and low estimates of benefits and costs.

A major barrier to efforts by local governments to adopt smoking bans is preemption, which is the passage or presence of a state law with weaker restrictions that prevents implementation and enforcement of stronger local laws. (Eliminating preemption statutes is one of the tobacco objectives of *Healthy People 2010*¹⁵ [Table 1-1].) Political opposition to legislative efforts to reduce ETS exposures in all workplaces and in a variety of public areas can be significant, and may include organizations representing tobacco users, businesses concerned about potential changes in revenue, and groups sponsored by the tobacco industry.

In conclusion, the Task Force recommends smoking bans and restrictions on the basis of strong evidence of effectiveness in decreasing both the amount of, and exposure to, environmental tobacco smoke. The findings of this review should be applicable to most indoor workers in the United States. Additional benefits of these interventions include reductions in daily consumption

of cigarettes among workers exposed to bans or restrictions and increases in tobacco use cessation by smokers exposed to workplace smoking bans.

Community Education to Reduce Exposure to Environmental Tobacco Smoke in the Home: Insufficient Evidence to Determine Effectiveness

Community education includes all efforts to increase knowledge and to change attitudes about the health effects of exposure to environmental tobacco smoke (ETS). Techniques may include mass media messages, small media messages (including educational materials), and counseling provided outside of healthcare settings.

Community education provides information to parents, other occupants, and visitors about the health risks for nonsmoking adults and for children caused by smoking in the home (infants and children get most of their ETS exposure in the home⁴³⁶). Information is disseminated to households to motivate (1) tobacco users to reduce exposures to ETS in the household by quitting or by smoking outdoors, (2) nonsmokers in the household to assist smokers in their efforts to quit, and (3) members of the household to establish home smoking bans or restrictions. The combination of reduced indoor smoking and increased cessation will result in a reduction in indoor ETS exposure and, therefore, a reduction in tobacco-related illness and death.

Effectiveness

- We found insufficient evidence to determine the effectiveness of community education to reduce exposure to ETS in the home.
- Evidence was insufficient because of the small number of available studies and limitations in their design and execution.
- Insufficient evidence means that we were not able to determine whether or not the intervention works.

The findings of our systematic review are based on one study.⁴³⁷ Two additional studies were identified but did not meet our quality criteria and were excluded from the review.^{438,439} The reviewed study evaluated the effect on infant exposure to ETS of education provided by nurses during home visits. At one-year follow-up, the difference in ETS exposure between intervention and comparison households was a reduction of 4 percentage points; overall exposure, however, increased in both study arms. The inconsistency of these findings provides insufficient evidence to determine the effectiveness of community education in reducing home exposure to ETS.

Because we could not establish the effectiveness of this intervention, we did not examine situations in which it would be applicable, information about economic efficiency, or possible barriers to implementation.

Potential benefits of education to reduce ETS exposure in the home include motivating tobacco users in the household to try to quit. No harms were identified.

In conclusion, the Task Force found insufficient evidence to determine the effectiveness of community education in reducing exposure to ETS in the home because of the small number of available studies and limitations in their design and execution.

USE OF THESE RECOMMENDATIONS

The evidence reviews and recommendations presented in this chapter provide a list of effective options to support efforts by states, communities, and healthcare systems to reduce tobacco use and exposures to environmental tobacco smoke. Prevention programs and planners can compare their current prevention activities with these recommendations, take steps to ensure that existing interventions are adequately implemented and funded, and consider additional interventions in the ongoing effort to build and maintain comprehensive tobacco prevention programs.

Based on evidence of effectiveness documented in the scientific literature, recommendations from the Task Force support the following population-based tobacco prevention and control efforts:

- Clean indoor air legislation prohibiting tobacco use in indoor public and private workplaces.
- Federal, state, and local efforts to increase tobacco product excise taxes as an effective public health intervention to promote tobacco use cessation and to reduce the initiation of tobacco use among youth.
- The funding and implementation of long-term, high-intensity mass media campaigns using paid broadcast times and media messages developed through formative research.
- Proactive telephone cessation support services (quit lines).
- Reduced or eliminated co-payments for effective cessation therapies.
- Reminder systems for healthcare providers.
- Combinations of efforts to mobilize communities to identify and reduce the commercial availability of tobacco products to youth.

In reflecting the available evidence on effectiveness, recommendations from the Task Force confirm the importance of coordinated or combined intervention efforts in tobacco prevention. Evidence of effectiveness in efforts to reduce tobacco use among youth through access restrictions, to disseminate anti-tobacco messages through mass media, and to assist tobacco users in their efforts to quit via telephone comes predominantly from studies that implemented these interventions in combination with other strategies.

The contribution to our available evidence on the effectiveness of multi-component intervention efforts should be fully appreciated by program planners. The available evidence on effectiveness for any single-component intervention is small and is concentrated in evaluations of policy interventions. In this review, only one single-component program intervention—healthcare provider reminder systems—had a sufficient body of evidence on effectiveness to support a Task Force recommendation for use. For the multicomponent reviews described in this chapter, readers are encouraged to go beyond the short recommendation summaries and to learn more about the intervention combinations described in the studies contributing to the recommendation from the Task Force.

A comprehensive tobacco prevention program is often interpreted to represent concurrent intervention efforts to promote or assist tobacco use cessation, to reduce tobacco use initiation, and to implement clean indoor air policies. The *Community Guide* reviews suggest that effectiveness within each of these strategic directions (cessation, initiation, reducing secondhand smoke) demands a similar comprehensive approach in the combination and coordination of interventions.

Recommendations from the Task Force also confirm the effectiveness (and the importance) of laws and policies in tobacco prevention. State and community clean indoor air laws are effective in reducing exposure to secondhand tobacco smoke. Excise taxes on tobacco products are effective in reducing tobacco use among both adults and youth. Healthcare system policies reducing out-of-pocket costs for effective cessation therapies increase and improve client cessation efforts. Although public policies have political implications and opposition, these effective intervention options should not be prematurely “taken off the table” for consideration by prevention programs and planners.

Although these reviews include summary measurements of the magnitude of effect, readers are strongly advised to also consider the size and scope of their target population for the intervention when drawing comparisons. Mass media campaigns and policies (excise tax increases, clean indoor air legislation) deliver an intervention impact across a very broad (and potentially very large) population. Interventions with effects of relatively small magnitude, when applied across an entire community, can contribute to significant change within the population.

In many states, preemption is a major barrier to the implementation of effective tobacco prevention policies and programs. It is a direct obstacle to the adoption of local clean indoor air ordinances and to local efforts to restrict youth access to tobacco products from commercial sources. In settings where effective options are blocked, programs should determine whether or not effective intervention combinations are still feasible to address the prevention

objective. In some cases, program resources may be better spent on implementing effective interventions to address other prevention goals. Programs must also remain vigilant for the introduction of new preemption legislation.

The work of the Task Force on Community Preventive Services complements an array of useful evidence reviews and guidelines^{2,3,6,17,84,440,441} available to tobacco prevention and control programs. These reviews, employing different methods to select, appraise, and summarize the available evidence, report similar findings and evidence-based conclusions.

In summary, *Community Guide* reviews provide programs and planners with a (1) concise summary on effectiveness of interventions and recommendations for use; (2) convenient guide to specific studies contributing to the evidence-based conclusions on effectiveness; and (3) complementary review to support similar conclusions across the available guidelines. The reviews and the recommendations provided in this chapter, therefore, can contribute in different ways to tobacco prevention and control efforts across a range of audiences, settings, and situations.

CONCLUSION

This chapter summarizes conclusions and recommendations to date from the Task Force on interventions to reduce the initiation of tobacco use, to increase tobacco use cessation, and to reduce exposure to ETS. To reduce the initiation of tobacco use, the Task Force recommends increasing the unit price for tobacco products; mass media education campaigns combined with other interventions; and community mobilization combined with additional interventions to restrict minors' access to tobacco products. Evidence was insufficient to determine the effectiveness of restricting minors' access to tobacco through the following interventions when implemented alone: sales laws directed at retailers; laws directed at minors' purchase, possession, or use of tobacco products; active enforcement of sales laws; retailer education with reinforcement; retailer education without reinforcement; and community education.

To increase cessation of tobacco use, the Task Force recommends increasing the unit price for tobacco products; mass media campaigns combined with other interventions; healthcare provider reminder systems; provider reminders combined with provider education; reducing client out-of-pocket costs for effective cessation therapies; and multicomponent interventions that include client telephone support (quit lines). Evidence was insufficient to determine the effectiveness of mass media cessation series; mass media cessation contests; healthcare provider education alone; and healthcare provider feedback and assessment.

To reduce exposure to ETS, the Task Force recommends smoking bans and

restrictions. Evidence was insufficient to determine the effectiveness of community education to reduce exposure to ETS in the home.

Details of these reviews have been published¹⁸⁻¹⁹ and these articles, along with additional information about the reviews, are available at www.thecommunityguide.org/tobacco.

Acknowledgments

This chapter was written by the members of the systematic review development team: Jonathan E. Fielding, MD, MPH, MBA, Los Angeles Department of Health Services, University of California Los Angeles School of Public Health, University of California Los Angeles School of Medicine, Los Angeles, California, and the Task Force on Community Preventive Services; Peter A. Briss, MD, Division of Prevention Research and Analytic Methods (DPRAM), Epidemiology Program Office (EPO), Centers for Disease Control and Prevention (CDC), Atlanta; Vilma G. Carande-Kulis, MS, PhD, DPRAM/EPO/CDC; David P. Hopkins, DPRAM/EPO/CDC; Corinne G. Husten, MD, MPH; Terry F. Pechacek, PhD, Office on Smoking and Health (OSH), National Centers for Chronic Disease Prevention and Health Promotion (NCCDPHP), Centers for Disease Control and Prevention, Atlanta; Tom J. Glynn, PhD, American Cancer Society, Washington, DC.

Consultants for all reviews were: Dileep G. Bal, MD, California Department of Health Services, Sacramento; Anthony Biglan, PhD, Oregon Research Institute, Eugene; Patricia A. Buffler, PhD, MPH, University of California, Berkeley and the Task Force on Community Preventive Services; Gregory Connolly, DMD, MPH, Massachusetts Tobacco Control Program, Boston; K. Michael Cummings, PhD, MPH, Roswell Park Institute, Buffalo, New York; Michael C. Fiore, MD, MPH, University of Wisconsin Medical School, Madison; David W. Fleming, MD, Centers for Disease Control and Prevention, Atlanta; Sally Malek, MPH, North Carolina Department of Health, Raleigh; Patricia A. Mullen, DrPH, University of Texas Health Sciences Center, Houston and the Task Force on Community Preventive Services; Cheryl L. Perry, PhD, University of Minnesota, Minneapolis; John P. Pierce, PhD, University of California, San Diego; Helen H. Schauffler, PhD, University of California, Berkeley; Randy H. Schwartz, MSPH, Maine Bureau of Health, Augusta; Mitchell Zeller, American Legacy Foundation, Washington, DC.

Consultants who worked on specific intervention reviews were: Namita Sharma MA, MPA; Trevor A. Woollery, PhD; Don J. Sharp, MD, DTMH; Jeff W. McKenna, MS; Corinne G. Husten, MD, MPH, OSH/NCCDPHP/CDC, Atlanta; Connie J. Ricard, MPH; Jeffrey R. Harris, MD, MPH, DPRAM/EPO/CDC, Atlanta.

Articles included in the reviews were abstracted by: J.F. Bond, MS; C.D. Archbald, MD, MPH; B.M. Morissette, MPH, PhD; J.A. Dake, MPH; C. Herrington, MPH; N.L. Lee, BS; L.M. Tomich, MS, RD; Y. Yousey, MS, RN.

References

1. McGinnis JM, Foege WH. Actual causes of death in the United States. *JAMA* 1993;270:2207-12.
2. U.S. Department of Health and Human Services. Reducing tobacco use: a report of

the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Office on Smoking and Health, 2000.

3. Centers for Disease Control and Prevention. Best practices for comprehensive tobacco control programs—August 1999. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1999.

4. Centers for Disease Control and Prevention. Cigarette smoking among adults—United States, 2002. *MMWR* 2004;53(20): 427–31.

5. U.S. Department of Health and Human Services. Tobacco use among U.S. racial/ethnic minority groups—African Americans, American Indians and Alaska Natives, Asian Americans and Pacific Islanders, and Hispanics: a report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1998.

6. Fiore MC, Bailey WC, Cohen SJ, et al. Treating tobacco use and dependence. Clinical practice guideline. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, 2000. Available at: <http://www.surgeongeneral.gov/tobacco>. Accessed July 13, 2000.

7. National Institutes of Health. Tobacco and the clinician: interventions for medical and dental practice. Washington, DC: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, 1994. NIH Pub No 94–3693.

8. U.S. Department of Health and Human Services. Reducing the health consequences of smoking: 25 years of progress. A report of the Surgeon General. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, 1989. DHHS Pub No (CDC) 89–8411.

9. U.S. Department of Health and Human Services. The health benefits of smoking cessation. A report of the Surgeon General. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1990. DHHS Pub No (CDC) 90–8416.

10. U.S. Department of Health and Human Services. Preventing tobacco use among young people: a report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1994.

11. University of Michigan, Institute for Social Research. Monitoring the Future Survey. Available at: <http://www.monitoringthefuture.org/data/03data.html#2003data-cigs>. Accessed March 16, 2004.

12. U.S. Environmental Protection Agency. Respiratory health effects of passive smoking: lung cancer and other disorders. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, 1992. EPA/600/6–90/006F.

13. California Environmental Protection Agency. Health effects of exposure to environmental tobacco smoke—final report and appendices. Sacramento: California Environmental Protection Agency, Office of Environmental Health Hazard Assessment, 1997.

14. Centers for Disease Control and Prevention. Second national report on human exposure to environmental chemicals. Available at: <http://www.cdc.gov/exposurereport/tobacco/pdf/tobaccosmoke.pdf>. Accessed March 16, 2004.

15. U.S. Department of Health and Human Services. *Healthy people 2010*. 2nd ed. Washington, DC: U.S. Government Printing Office, 2000.

16. Hopkins DP, Husten CG, Fielding JE, Rosenquist JN, Westphal LL. Evidence reviews and recommendations on interventions to reduce tobacco use and exposure to environmental tobacco smoke. A summary of selected guidelines. *Am J Prev Med* 2001; 20(2S):67–87.

17. U.S. Preventive Services Task Force. *Guide to clinical preventive services: report of the U.S. Preventive Services Task Force*. 2nd ed. Baltimore: Williams & Wilkins, 1996.

18. Centers for Disease Control and Prevention. *Strategies for reducing exposure to environmental tobacco smoke, increasing tobacco-use cessation, and reducing initiation in communities and health-care systems. A report on recommendations of the Task Force on Community Preventive Services*. *MMWR* 2000;49(No. RR-12):1–11.

19. Task Force on Community Preventive Services. Recommendations regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Prev Med* 2001;20(2S):10–5.

20. Hopkins DP, Briss PA, Ricard CJ, et al. Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Prev Med* 2001;20(2S):16–66.

21. Lynch BS, Bonnie RJ, Institute of Medicine: Committee on Preventing Nicotine Addiction in Children and Youths. *Overview: growing up tobacco free. Preventing nicotine addiction in children and youths*. Washington, DC: National Academy Press, 1994.

22. Chaloupka F, Warner K. The economics of smoking. In: Culyer AJ, Newhouse JP, eds. *Handbook of health economics*. Vol. 1B. Amsterdam: Elsevier Science, 2000: 1539–627.

23. Centers for Disease Control and Prevention. Response to increases in cigarette prices by race/ethnicity, income, and age groups—United States, 1976–1993. *MMWR* 1998;47(29):605–9.

24. Chaloupka FJ, Pacula RL. Sex and race differences in young people's responsiveness to price and tobacco control policies. *Tob Control* 1999;8(4):373–7.

25. Chaloupka FJ, Wechsler H. Price, tobacco control policies and smoking among young adults. *J Health Econ* 1997;16:359–73.

26. Gruber J. Youth smoking in the U.S.: prices and policies. Available at: <http://papers.nber.org/papers/w7506>. Accessed April 12, 2004.

27. Lewit EM, Coate D, Grossman M. The effects of government regulation on teenage smoking. *J Polit Econ* 1981;24:545–70.

28. Lewit EM, Hyland A, Kerrebrock N, Cummings KM. Price, public policy, and smoking in young people. *Tob Control* 1997;6(suppl 2):S17–S24.

29. Ohsfeldt RL, Boyle RG, Capilouto E. Effects of tobacco excise taxes on the use of smokeless tobacco products in the USA. *Health Educ* 1997;6:525–31.

30. Wasserman J, Manning WG, Newhouse JP, Winkler JD. The effects of excise taxes and regulations on cigarette smoking. *J Health Econ* 1991;10:43–64.

31. Chaloupka FJ, Grossman M. Price, tobacco control policies and youth smoking. *National Bureau of Economic Research* 1996;Working Paper 5740:1–39.

32. Chaloupka FJ, Tauras JA, Grossman M. Public policy and youth smokeless tobacco use. *South Econ J* 1997;64(2):503–16.

33. Ohsfeldt RL, Boyle RG. Tobacco excise taxes and rates of smokeless tobacco use in the U.S.: an exploratory ecological analysis. *Tob Control* 1994;3:316–23.

34. Koh HK. An analysis of the successful 1992 Massachusetts tobacco tax initiative. *Tob Control* 1996;5(3):220–5.
35. Nicholl J. Tobacco tax initiatives to prevent tobacco use. *Cancer* 1998;83:2666–79.
36. Traynor MP, Glantz SA. California's tobacco tax initiative: the development and passage of Proposition 99. *J Health Polit Policy Law* 1996;21(3):543–85.
37. Worden JK, Flynn BS. Effective use of mass media to prevent cigarette smoking. *J Public Health Manag Pract* 2000;6(3):vii–viii.
38. Bauman KE, LaPrelle J, Brown JD, Koch GG, Padgett CA. The influence of three mass media campaigns on variables related to adolescent cigarette smoking: results of a field experiment. *Am J Public Health* 1991;81(5):597–604.
39. Centers for Disease Control and Prevention. Tobacco use among middle and high school students—Florida, 1998 and 1999. *MMWR* 1999;48(12):248–53.
40. Flay BR, Hansen WB, Johnson CA, et al. Implementation effectiveness trial of a social influences smoking prevention program using schools and television. *Health Educ Res* 1987;2(4):385–400.
41. Flay BR, Miller TQ, Hedeker D, et al. The television, school, and family smoking prevention and cessation project. VIII. Student outcomes and mediating variables. *Prev Med* 1995;24(1):29–40.
42. Flynn BS, Worden JK, Secker-Walker RH, Pirie PL, Badger GJ, Carpenter JH. Long-term responses of higher and lower risk youths to smoking prevention interventions. *Prev Med* 1997;26(3):389–94.
43. Hafstad A, Straypedersen B, Langmark F. Use of provocative emotional appeals in a mass media campaign designed to prevent smoking among adolescents. *Eur J Public Health* 1997;7(2):122–7.
44. Johnson CA, Pentz MA, Weber MD, et al. Relative effectiveness of comprehensive community programming for drug abuse prevention with high-risk and low-risk adolescents. *J Consult Clin Psychol* 1990;58(4):447–56.
45. Kaufman JS, Jason LA, Sawlski LM, Halpert JA. A comprehensive multi-media program to prevent smoking among black students. *J Drug Educ* 1994;24(2):95–108.
46. Murray DM, Prokhorov AV, Harty KC. Effects of a statewide antismoking campaign on mass media messages and smoking beliefs. *Prev Med* 1994;23(1):54–60.
47. Perry CL, Kelder SH, Murray DM, Klepp KI. Communitywide smoking prevention: long-term outcomes of the Minnesota Heart Health Program and the Class of 1989 Study. *Am J Public Health* 1992;82(9):1210–6.
48. Siegel M, Biener L. The impact of an antismoking media campaign on progressions to established smoking: results of a longitudinal youth study. *Am J Public Health* 2000;90(3):1–7.
49. Vartiainen E, Paavola M, McAlister A, Puska P. Fifteen-year follow-up of smoking prevention effects in the North Karelia Youth Project. *Am J Public Health* 1998;88(1):81–5.
50. Meister JS. Designing an effective statewide tobacco control program—Arizona. *Cancer* 1998;83(12 suppl rep):2728–32.
51. Seghers T, Foland S. Anti-tobacco media campaign for young people. *Tob Control* 1998;7(suppl):S29–S30 (discussion).
52. Bauman KE, Brown JD, Bryan ES, Fisher LA, Padgett CA, Sweeney JM. Three mass media campaigns to prevent adolescent cigarette smoking. *Prev Med* 1988;17(5):510–30.

53. Bauman KE, Padgett CA, Koch GG. A media-based campaign to encourage personal communication among adolescents about not smoking cigarettes: participation, selection and consequences. Special issue: smoking. *Health Educ Res* 1989;4(1):35–44.
54. Brannon BR, Dent CW, Flay BR, et al. The television, school, and family project. V. The impact of curriculum delivery format on program acceptance. *Prev Med* 1989;18:492–502.
55. Flay BR, Brannon BR, Johnson CA, et al. The television school and family smoking prevention and cessation project. 1. Theoretical basis and program development. *Prev Med* 1988;17(5):585–607.
56. Florida Department of Health. Florida Tobacco Youth Survey. Available at: http://www.doh.state.fl.us/disease_ctrl/epi/FYTS/index.htm. Accessed March 18, 2004.
57. Flynn BS, Worden JK, Secker-Walker RH, Badger GJ, Geller BM, Costanza MC. Prevention of cigarette smoking through mass media intervention and school programs. *Am J Public Health* 1992;82(6):827–34.
58. Flynn BS, Worden JK, Secker-Walker RH, et al. Mass media and school interventions for cigarette smoking prevention: effects 2 years after completion. *Am J Public Health* 1994;84(7):1148–50.
59. Hafstad A, Aaro LE, Langmark F. Evaluation of an anti-smoking mass media campaign targeting adolescents: the role of affective responses and interpersonal communication. *Health Educ Res* 1996;11(1):29–38.
60. Hafstad A, Aaro LE. Activating interpersonal influence through provocative appeals—evaluation of a mass media-based antismoking campaign targeting adolescents. *Health Commun* 1997;9(3):253–72.
61. Hafstad A, Aaro LE, Engeland A, Andersen A, Langmark F, Straypedersen B. Provocative appeals in anti-smoking mass media campaigns targeting adolescents—the accumulated effect of multiple exposures. *Health Educ Res* 1997;12(2):227–36.
62. MacKinnon DP, Johnson CA, Pentz MA, et al. Mediating mechanisms in a school-based drug prevention program: first-year effects of the Midwestern Prevention Project. *Health Psychol* 1991;10(3):164–72.
63. Murray DM, Perry CL, Griffin G, et al. Results from a statewide approach to adolescent tobacco use prevention. *Prev Med* 1992;21(4):449–72.
64. Pentz MA, MacKinnon DP, Flay BR, Hansen WB, Johnson CA, Dwyer JH. Primary prevention of chronic diseases in adolescence: effects of the Midwestern Prevention Project on tobacco use. *Am J Epidemiol* 1989;130(4):713–24.
65. Pentz MA, MacKinnon DP, Dwyer JH, et al. Longitudinal effects of the Midwestern Prevention Project on regular and experimental smoking in adolescents. *Prev Med* 1989;18(2):304–21.
66. Pentz MA, Johnson A, Dwyer JH, MacKinnon DM, Hansen WB, Flay BR. A comprehensive community approach to adolescent drug abuse prevention: effects on cardiovascular disease risk behaviors. *Ann Med* 1989;21(3):219–22.
67. Pentz MA, Trebow EA, Hansen WB, et al. Effects of program implementation on adolescent drug use behavior: the Midwestern Prevention Project (MPP). *Eval Rev* 1990;14(3):264–89.
68. Perry CL, Klepp KI, Sillers C. Community-wide strategies for cardiovascular health: the Minnesota Heart Health Program youth program. Special issue: smoking. *Health Educ Res* 1989;4(1):87–101.
69. Puska P, Vartiainen E, Pallonen U, et al. The North Karelia Youth Project. A

community-based intervention study on CVD risk factors among 13- to 15-year-old children: study design and preliminary findings. *Prev Med* 1981;10(2):133–48.

70. Puska P, Vartiainen E, Pallonen U, et al. The North Karelia Youth Project: evaluation of two years of intervention on health behavior and CVD risk factors among 13- to 15-year old children. *Prev Med* 1982;11(5):550–70.

71. Secker-Walker RH, Worden JK, Holland RR, Flynn BS, Detsky AS. A mass media programme to prevent smoking among adolescents: costs and cost effectiveness. *Tob Control* 1997;6:207–12.

72. Sussman S, Brannon BR, Flay BR, et al. The television, school, and family smoking prevention/cessation project. II. Formative evaluation of television segments by teenagers and parents—implications for parental involvement in drug education. *Health Educ Res* 1986;1(3):185–94.

73. Sussman S, Dent CW, Brannon BR, et al. The television, school and family smoking prevention/cessation project. IV. Controlling for program success expectancies across experimental and control conditions. *Addict Behav* 1989;14(6):601–10.

74. Vartiainen E, Pallonen U, McAlister A, Koskela K, Puska P. Effect of two years of educational intervention on adolescent smoking (the North Karelia Youth Project). *Bull World Health Organ* 1983;61(3):529–32.

75. Vartiainen E, Pallonen U, McAlister A, Koskela K, Puska P. Four-year follow-up results of the smoking prevention program in the North Karelia Youth Project. *Prev Med* 1986;15(6):692–8.

76. Vartiainen E, Fallonen U, McAlister AL, Puska P. Eight-year follow-up results of an adolescent smoking prevention program: the North Karelia Youth Project. *Am J Public Health* 1990;80(1):78–9.

77. Vartiainen E, Tossavainen K, Viri L, Niskanen E, Puska P. The North Karelia Youth Programs. *Ann N Y Acad Sci* 1991;623:332–49.

78. Worden JK, Flynn BS, Geller BM, et al. Development of a smoking prevention mass media program using diagnostic and formative research. *Prev Med* 1988;17(5):531–58.

79. Worden JK, Flynn BS, Solomon LJ, Secker-Walker RH, Badger GJ, Carpenter JH. Using mass media to prevent cigarette smoking among adolescent girls. *Health Educ Q* 1996;23(4):453–68.

80. Forster JL, Wolfson M. Youth access to tobacco: policies and politics. *Annu Rev Public Health* 1998;19:203–35.

81. Forster JL, Wolfson M, Murray DM, Wagenaar AC, Claxton AJ. Perceived and measured availability of tobacco to youths in 14 Minnesota communities: the TPOP Study. *Tobacco Policy Options for Prevention*. *Am J Prev Med* 1997;13(3):167–74.

82. Harrison PA, Fulkerson JA, Park E. The relative importance of social versus commercial sources in youth access to tobacco, alcohol, and other drugs. *Prev Med* 2000;31(1):39–48.

83. Hinds MW. Impact of a local ordinance banning tobacco sales to minors. *Public Health Rep* 1992;107(3):355–8.

84. Institute of Medicine. Growing up tobacco free: preventing nicotine addiction in children and youths. Lynch BS, Bonnie RJ, eds. Washington, DC: National Academy Press, 1994.

85. Altman DG, Rasenick-Douss L, Foster V, Tye JB. Sustained effects of an educational program to reduce sales of cigarettes to minors. *Am J Public Health* 1991;81(7):891–3.

86. Altman DG, Wheelis AY, McFarlane M, Lee H, Fortmann SP. The relationship between tobacco access and use among adolescents: a four community study. *Soc Sci Med* 1999;48(6):759–75.
87. Biglan A, Ary DV, Smolkowski K, Duncan T, Black C. A randomised controlled trial of a community intervention to prevent adolescent tobacco use. *Tob Control* 2000; 9(1):24–32.
88. Chapman S, King M, Andrews B, McKay E, Markham P, Woodward S. Effects of publicity and a warning letter on illegal cigarette sales to minors. *Aust J Public Health* 1994;18(1):39–42.
89. Feighery E, Altman DG, Shaffer G. The effects of combining education and enforcement to reduce tobacco sales to minors. A study of four northern California communities. *JAMA* 1991;266(22):3168–71.
90. Forster JL, Murray DM, Wolfson M, Blaine TM, Wagenaar AC, Hennrikus DJ. The effects of community policies to reduce youth access to tobacco. *Am J Public Health* 1998;88(8):1193–8.
91. Jason LA, Billows WD, Schnopp-Wyatt DL, King C. Long term findings from Woodridge in reducing illegal cigarette sale to older minors. *Eval Health Prof* 1996;19(1):3–13.
92. Wildey MB, Woodruff SI, Pampalone SZ, Conway TL. Self-service sale of tobacco: how it contributes to youth access. *Tob Control* 1995;4:355–61.
93. Junck E, Humphries J, Rissel C. Reducing tobacco sales to minors in Manly: 10 months follow-up. *Health Promot J Aust* 1997;70:29–34.
94. Altman DG, Foster V, Rasenick-Douss L, Tye JB. Reducing the illegal sale of cigarettes to minors. *JAMA* 1989;261(1):80–3.
95. Biglan A, Ary DV, Koehn V, et al. Mobilizing positive reinforcement in communities to reduce youth access to tobacco. *Am J Community Psychol* 1996;24(5):625–38.
96. Jason LA, Ji PY, Anes MD, Birkhead SH. Active enforcement of cigarette control laws in the prevention of cigarette sales to minors [see comments]. *JAMA* 1991;266(22): 3159–61.
97. Keay KD, Woodruff SI, Wildey MB, Kenney EM. Effect of retailer intervention on cigarette sales to minors in San Diego County, California. *Tob Control* 1993;2:145–51.
98. Biglan A, Henderson J, Humphrey D. Mobilising positive reinforcement to reduce youth access to tobacco. *Tob Control* 1995;4:42–8.
99. Erickson AD, Woodruff SI, Wildey MB, Kenney E. A baseline assessment of cigarette sales to minors in San Diego, California. *J Community Health* 1993;18(4):213–24.
100. Jason LA, Katz R, Vavra J, Schnopp-Wyatt DL, Talbot B. Long-term follow-up of youth access to tobacco laws' impact on smoking prevalence. *J Hum Behav Soc Environ* 1999;2(3):1–13.
101. Cummings KM, Coogan K. Organizing communities to prevent the sale of tobacco products to minors. *Int Q Community Health Educ* 1992;13(1):77–86.
102. Dovell RA, Mowat DL, Dorland J, Lam M. Changes among retailers selling cigarettes to minors. *Can J Public Health* 1996;87(1):66–8.
103. Lewis RK, Paine-Andrews A, Fawcett SB, et al. Evaluating the effects of a community coalition's efforts to reduce illegal sales of alcohol and tobacco products to minors. *J Community Health* 1996;21(6):429–36.
104. Schensky AE, Smith SS, Icenogle DL, Fiore MC. Youth tobacco sale compliance checks: impact on vendor practices and community policy. *Wis Med J* 1996;95(11): 775–8.

105. Staff M, March L, Brnabic A, et al. Can non-prosecutory enforcement of public health legislation reduce smoking among high school students? [see comments]. *Aust N Z J Public Health* 1998;22(3 suppl):332-5.

106. Centers for Disease Control and Prevention. Illegal sales of cigarettes to minors—Ciudad Juarez, Mexico; El Paso, Texas; and Las Cruces, New Mexico, 1999. *MMWR* 1999;48:394-8.

107. Naidoo J, Platts C. Smoking prevention in Bristol: getting maximum results using minimum resources. *Health Educ J* 1985;44:39-42.

108. Rigotti NA, DiFranza JR, Chang Y, Tisdale T, Kemp B, Singer DE. The effect of enforcing tobacco-sales laws on adolescents' access to tobacco and smoking behavior. *N Engl J Med* 1997;337(15):1044-51.

109. Alcaraz R, Klonoff EA, Landrine H. The effects on children of participating in studies of minors' access to tobacco. *Prev Med* 1997;26(2):236-40.

110. DiFranza JR, Peck RM, Radecki TE, Savageau JA. What is the potential cost-effectiveness of enforcing a prohibition on the sale of tobacco to minors? *Prev Med* 2001;32(2):168-74.

111. Siegel M, Carol J, Jordan J, et al. Preemption in tobacco control: review of an emerging public health problem. *JAMA* 1997;278:858-63.

112. Gorovitz E, Mosher J, Pertschuk M. Preemption or prevention?: Lessons learned from efforts to control firearms, alcohol, and tobacco. *J Public Health Policy* 1998;19(1):36-50.

113. Jacobson PD, Wasserman J. The implementation and enforcement of tobacco control laws: policy implications for the activists and the industry. *J Health Polit Policy Law* 1999;24(3):567-98.

114. Mosher JF. The merchants, not the customers: resisting the alcohol and tobacco industries' strategy to blame young people for illegal alcohol and tobacco sales. *J Public Health Policy* 1995;16(4):412-32.

115. DiFranza JR. The empire strikes back. *Tob Access Law News* 1994;25:1-8.

116. DiFranza JR, Godshall B. Tobacco industry lobbies to sabotage Synar law. *Tobacco-Free Youth Rep* 1994;6(3):7-10.

117. Bidell MP, Furlong MJ, Dunn DM, Koegler JE. Case study of attempts to enact self service tobacco display ordinances: a tale of three communities. *Tob Control* 2000;9:71-7.

118. Forster JL, Hourigan ME, Kelder S. Locking devices on cigarette vending machines: evaluation of a city ordinance. *Am J Public Health* 1992;82(9):1217-9.

119. Siegel M, Biener L, Rigotti NA. The effect of local tobacco sales laws on adolescent smoking initiation. *Prev Med* 1999;29(5):334-42.

120. Cismoski J. Blinded by the light: the folly of tobacco possession laws against minors. *Wis Med J* 1994;93(11):591-8.

121. Lazovich D, Ford J, Forster J, Riley B. A pilot study to evaluate a tobacco diversion program. *Am J Public Health* 2001;91(11):1790-1.

122. Gemson DH, Moats HL, Watkins BX, Ganz ML, Robinson S, Heaton E. Laying down the law: reducing illegal tobacco sales to minors in Central Harlem. *Am J Public Health* 1998;88(6):936-9.

123. Bagott M, Jordan C, Wright C, Jarvis S. How easy is it for young people to obtain cigarettes, and do test sales by trading standards have any effect? A survey of two schools in Gateshead. *Child Care Health Dev* 1998;24(3):207-16.

124. McDermott SR, Scott KL, Frintner MP. Accessibility of cigarettes to minors in suburban Cook County, Illinois. *J Community Health* 1998;23(2):153–60.
125. Woodruff SI, Wildey MB, Conway TL, Clapp EJ. Effect of a brief retailer intervention to reduce the sale of single cigarettes. *Am J Health Promot* 1995;9(3):172–4.
126. Skretny MT, Cummings KM, Sciandra R, Marshall J. An intervention to reduce the sale of cigarettes to minors. *N Y State J Med* 1990;90(2):54–5.
127. Schofield MJ, Sanson-Fisher RW, Gulliver S. Interventions with retailers to reduce cigarette sales to minors: a randomised controlled trial. *Aust N Z J Public Health* 1997;21(6):590–6.
128. Voorhees CC, Yanek LR, Stillman FA, Becker DM. Reducing cigarette sales to minors in an urban setting: issues and opportunities for merchant intervention. *Am J Prev Med* 1998;14(2):138–42.
129. Abernathy TJ. Compliance for Kids: a community-based tobacco prevention project. *Can J Public Health* 1994;85(2):82–4.
130. DiFranza JR, Brown LJ. The Tobacco Institute's "It's the Law" campaign: has it halted illegal sales of tobacco to children? *Am J Public Health* 1992;82(9):1271–3.
131. National Cancer Institute. Population-based smoking cessation: proceedings of a conference on what works to influence cessation in the general population. Smoking and tobacco control monograph No. 12. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, 2000. NIH Pub. No 00–4892.
132. Flewelling RL, Kenney E, Elder JP, Pierce J, Johnson M, Bal DG. First-year impact of the 1989 California cigarette tax increase on cigarette consumption. *Am J Public Health* 1992;82(6):867–9.
133. Glantz S. Changes in cigarette consumption, prices, and tobacco industry revenues associated with California's Proposition 99. *Tob Control* 1993;2:311–4.
134. Hu TW, Bai J, Keeler TE, Barnett PG, Sung HY. The impact of California Proposition 99, a major anti-smoking law, on cigarette consumption. *J Public Health Policy* 1994;15(1):26–36.
135. Hu TW, Sung HY, Keeler TE. Reducing cigarette consumption in California: tobacco taxes vs an anti-smoking media campaign. *Am J Public Health* 1995;85(9):1218–22.
136. Hu TW, Ren QF, Keeler TE, Bartlett J. The demand for cigarettes in California and behavioural risk factors. *Health Econ* 1995;4(1):7–14.
137. Keeler TE, Hu T, Barnett PG, Manning WG. Taxation, regulation, and addiction: a demand function for cigarettes based on time-series evidence. *J Health Econ* 1993;12:1–18.
138. Pierce JP, Gilpin E, Emery SL, White MM, Rosbrook B, Berry C. Has the California Tobacco Control Program reduced smoking? *JAMA* 1998;280(10):893–9.
139. Centers for Disease Control and Prevention. Cigarette smoking before and after an excise tax increase and an antismoking campaign—Massachusetts, 1990–1996. *MMWR* 1996;45(44):966–70.
140. Centers for Disease Control and Prevention. Decline in cigarette consumption following implementation of a comprehensive tobacco prevention and education program—Oregon, 1996–1998. *MMWR* 1999;48(7):140–3.
141. Sung H, Hu T, Keeler TE. Cigarette taxation and demand: an empirical model. *Contemp Econ Policy* 1994;12:91–100.

142. Meier KJ, Licari MJ. The effect of cigarette taxes on cigarette consumption, 1955 through 1994. *Am J Public Health* 1997;87(7):1126–30.

143. Evans WN, Ringel JS, Stech D. Tobacco taxes and public policy to discourage smoking. In: Tax policy and the economy—Vol. 13. Poterba JM, ed. Cambridge, MA: MIT Press, 1999:1–56.

144. Baltagi BH, Goel RK. Quasi-experimental price elasticities of cigarette demand and the bootlegging effect. *Am Agric Econ Assoc* 1987;69(4):750–4.

145. Barnett PG, Keeler TE, Hu T. Oligopoly structure and the incidence of cigarette excise taxes. *J Public Econ* 1995;57:457–70.

146. Becker GS, Grossman M, Murphy KM. An empirical analysis of cigarette addiction. *Am Econ Rev* 1994;84(3):396–418.

147. Chaloupka F, Saffer H. Clean indoor air laws and the demand for cigarettes. *Contemp Policy Issues* 1992;10:72–83.

148. Evans WN, Farrelly MC. The compensating behavior of smokers: taxes, tar, and nicotine. *Rand J Econ* 1998;29(3):578–95.

149. Jackson JD, Saba RP. Some limits on taxing sin: cigarette taxation and health care finance. *South Econ J* 1997;63:761–75.

150. Moore MJ. Death and tobacco taxes. *Rand J Econ* 1996;27(2):415–28.

151. Peterson DE, Zeger SL, Remington PL, Anderson HA. The effect of state cigarette tax increases on cigarette sales, 1955 to 1988. *Am J Public Health* 1992;82(1):94–6.

152. Saba RP, Beard TR, Ekelund RB, Ressler RW. The demand for cigarette smuggling. *Econ Inq* 1995;23:189–202.

153. Seldon BJ, Doroodian K. A simultaneous model of cigarette advertising: effects on demand and industry response to public policy. *Rev Econ Stat* 1989;71(4):673–7.

154. Seldon BJ, Boyd R. The stability of cigarette demand. *Appl Econ* 1991;23(2):319–26.

155. Tegene A. Kalman filter and the demand for cigarettes. *Appl Econ* 1991;23(7):1175–82.

156. Thursby JG, Thursby MC. Interstate cigarette bootlegging: extent, revenue losses, and effects of federal intervention. *National Tax J* 2000;53(1):59–77.

157. Tremblay CH, Tremblay VJ. The impact of cigarette advertising on consumer surplus, profit, and social welfare. *Contemp Econ Policy* 1995;13:113–24.

158. Goel RK, Morey MJ. The interdependence of cigarette and liquor demand. *South Econ J* 1995;62(2):451–9.

159. Baltagi BH, Levin D. Estimating dynamic demand for cigarettes using panel data: the effects of bootlegging, taxation and advertising reconsidered. *Rev Econ Stat* 1986;68(1):148–55.

160. Bishop JA, Yoo JH. Health scare, excise taxes and advertising ban in the cigarette demand and supply. *South Econ J* 1985;52(2):402–11.

161. Chaloupka FJ. Clean indoor air laws, addiction, and cigarette smoking. *Appl Econ* 1992;24:193–205.

162. Chaloupka FJ. Rational addictive behavior and cigarette smoking. *J Polit Econ* 1991;99(41):722–42.

163. Douglas S, Hariharan G. The hazard of starting smoking: estimates from a split population duration model. *J Health Econ* 1994;13:213–30.

164. Fujii ET. The demand for cigarettes: further empirical evidence and its implications for public policy. *Appl Econ* 1980;12:479–89.

165. Lewit EM, Coate D. The potential for using excise taxes to reduce smoking. *J Health Econ* 1982;1(2):121–45.
166. Schneider L, Klein B, Murphy KM. Governmental regulation of cigarette health information. *J Law Econ* 1981;24:575–612.
167. Young T. The demand for cigarettes: alternative specifications of Fujii's model. *Appl Econ* 1983;15:203–11.
168. Galbraith JW, Kaiserman M. Taxation, smuggling and demand for cigarettes in Canada: evidence from time-series data. *J Health Econ* 1997;16:287–301.
169. Hamilton VH, Levinton C, St-Pierre Y, Grimard F. The effect of tobacco tax cuts on cigarette smoking in Canada. *Can Med Assoc J* 1997;156(2):187–91.
170. Mummery WK, Hagen LC. Tobacco pricing, taxation, consumption and revenue: Alberta 1985–1995. *Can J Public Health* 1996;87(5):314–6.
171. Jones A. UK demand for cigarettes 1954–1986, a double-hurdle approach. *J Health Econ* 1989;8(1):133–41.
172. Townsend J, Roderick P, Cooper J. Cigarette smoking by socioeconomic group, sex, and age: effects of price, income, and health publicity. *BMJ* 1994;309(6959):923–7.
173. Townsend JL. Cigarette tax, economic welfare, and social class patterns of smoking. *Appl Econ* 1987;19:355–65.
174. Worgotter GF, Kunze M. Cigarette prices and cigarette consumption in Austria, 1955–1983. *N Y State J Med* 1986;86(9):478–9.
175. Pekurinen M. The demand for tobacco products in Finland. *Brit J Addict* 1989;84(10):1183–92.
176. Leu R. The effects of cigarette price and anti-smoking publicity on cigarette consumption in Switzerland. *Rev Epidemiol Sante Publique* 1979;27(4):359–62.
177. Leu R. Anti-smoking publicity, taxation, and the demand for cigarettes. *J Health Econ* 1984;3:101–16.
178. Laugesen M, Meads C. Advertising, price, income and publicity effects on weekly cigarette sales in New Zealand supermarkets. *Brit J Addict* 1991;86(1):83–9.
179. Kaiserman MJ, Rogers B. Tobacco consumption declining faster in Canada than in the U.S. *Am J Public Health* 1991;81(7):902–4.
180. Labeaga JM. Individual behaviour and tobacco consumption: a panel data approach. *Health Econ* 1993;2(2):103–12.
181. Laugesen M, Meads C. Tobacco advertising restrictions, price, income and tobacco consumption in OECD countries, 1960–1986. *Brit J Addict* 1991;86(10):1343–54.
182. Miller E. Taxes, cigarettes and the health of Maine citizens. *J Maine Med Assoc* 1977;68(2):58–62.
183. Reekie WD. Consumers' surplus and the demand for cigarettes. *Manage Decis Econ* 1994;15:223–34.
184. Advisory Commission on Intergovernmental Relations. Cigarette tax evasion: a second look. Washington, DC: Advisory Commission on Intergovernmental Relations, 1985.
185. Cummings KM, Sciandra R, Davis S, Rimer BK. Results of an antismoking media campaign utilizing the Cancer Information Service. *J Natl Cancer Inst Monogr* 1993; (14):113–8.
186. Fortmann SP, Taylor CB, Flora JA, Jatulis DE. Changes in adult cigarette smoking prevalence after 5 years of community health education: the Stanford Five-City Project. *Am J Epidemiol* 1993;137(1):82–96.

187. Jenkins CN, McPhee SJ, Le A, Pham GQ, Ha NT, Stewart S. The effectiveness of a media-led intervention to reduce smoking among Vietnamese-American men. *Am J Public Health* 1997;87(6):1031–4.
188. Ledwith F. Immediate and delayed effects of postal advice on stopping smoking. *Health Bull* 1984;42(6):332–44.
189. Luepker RV, Murray DM, Jacobs DJ, et al. Community education for cardiovascular disease prevention: risk factor changes in the Minnesota Heart Health Program. *Am J Public Health* 1994;84(9):1383–93.
190. Marin G, Perez-Stable EJ. Effectiveness of disseminating culturally appropriate smoking-cessation information: Programa Latino para Dejar de Fumar. *J Natl Cancer Inst Monogr* 1995;(18):155–63.
191. McAlister AL, Ramirez AG, Amezcua C, Pulley L, Stern MP, Mercado S. Smoking cessation in Texas-Mexico border communities: a quasi-experimental panel study. *Am J Health Promot* 1992;6(4):274–9.
192. McPhee SJ, Jenkins CN, Wong C, et al. Smoking cessation intervention among Vietnamese Americans: a controlled trial. *Tob Control* 1997;4(suppl 1):S16–S24.
193. Meyer AJ. Skills training in a cardiovascular health education campaign. *J Consult Clin Psychol* 1980;48(2):129–42.
194. Mudde AN, de Vries H, Dolders MG. Evaluation of a Dutch community-based smoking cessation intervention. *Prev Med* 1995;24(1):61–70.
195. Pierce JP, Macaskill P, Hill D. Long-term effectiveness of mass media led anti-smoking campaigns in Australia. *Am J Public Health* 1990;80(5):565–9.
196. Vartiainen E, Puska P, Jousilahti P, Korhonen HJ, Tuomilehto J, Nissinen A. Twenty-year trends in coronary risk factors in north Karelia and in other areas of Finland. *Int J Epidemiol* 1994;23(3):495–504.
197. Dwyer T, Pierce JP, Hannam CD, Burke N. Evaluation of the Sydney “Quit For Life” anti-smoking campaign. Part 2. Changes in smoking prevalence. *Med J Aust* 1986;144(7):344–7.
198. Farquhar JW, Maccoby N, Wood PD, et al. Community education for cardiovascular health. *Lancet* 1977;1(8023):1192–5.
199. Farquhar JW, Fortmann SP, Maccoby N, et al. The Stanford Five-City Project: design and methods. *Am J Epidemiol* 1985;122(2):323–34.
200. Farquhar JW, Fortmann SP, Flora JA, et al. Effects of communitywide education on cardiovascular disease risk factors. The Stanford Five-City Project. *JAMA* 1990;264(3):359–65.
201. Jacobs DR Jr, Luepker RV, Mittelmark MB, et al. Community-wide prevention strategies: evaluation design of the Minnesota Heart Health Program. *J Chronic Dis* 1986;39(10):775–88.
202. Marin BV, Perez-Stable EJ, Marin G, Hauck WW. Effects of a community intervention to change smoking behavior among Hispanics. *Am J Prev Med* 1994;10(6):340–7.
203. Marin G, Marin BV, Perez-Stable EJ, Sabogal F, Otero-Sabogal R. Changes in information as a function of a culturally appropriate smoking cessation community intervention for Hispanics. *Am J Community Psychol* 1990;18(6):847–64.
204. Perez-Stable EJ, Marin BV, Marin G. A comprehensive smoking cessation program for the San Francisco Bay area Latino community: Programa Latino para Dejar de Fumar. *Am J Health Promot* 1993;7(6):430–42.

205. Pierce JP, Dwyer T, Frape G, Chapman S, Chamberlain A, Burke N. Evaluation of the Sydney "Quit For Life" anti-smoking campaign. Part 1. Achievement of intermediate goals. *Med J Aust* 1986;144(7):341-4.
206. Pierce JP, Burns DM, Berry C, et al. Reducing tobacco consumption in California: Proposition 99 seems to work [letter]. *JAMA* 1991;265(10):1257-8.
207. Popham WJ, Potter LD, Hetrick MA, Muthen LK, Duerr JM, Johnson MD. Effectiveness of the California 1990-1991 tobacco education media campaign. *Am J Prev Med* 1994;10(6):319-26.
208. Puska P, Wiio J, McAlister A, et al. Planned use of mass media in national health promotion: the "Keys to Health" TV program in 1982 in Finland. *Can J Public Health* 1985;76(5):336-42.
209. Boyd NR, Sutton C, Orleans CT, et al. Quit Today: a targeted communications campaign to increase use of the cancer information service by African American smokers. *Prev Med* 1998;27(5 (Pt 2)):S50-S60.
210. Ossip-Klein DJ, Shapiro RM, Stiggins J. Freedom line: increasing utilization of a telephone support service for ex-smokers. *Addict Behav* 1984;9(2):227-30.
211. Pierce JP, Anderson DM, Romano RM, Meissner HI, Odenkirchen JC. Promoting smoking cessation in the United States: effect of public service announcements on the Cancer Information Service telephone line. *J Natl Cancer Inst* 1992;84(9):677-83.
212. Barber JJ, Grichting WL. Australia's media campaign against drug abuse. *Int J Addict* 1990;25(6):693-708.
213. Campion P, Owen L, McNeill A, McGuire C. Evaluation of a mass media campaign on smoking and pregnancy. *Addiction* 1994;89(10):1245-54.
214. Gredler B, Kunze M. Impact of a national campaign on smoking attitudes and patterns in Austria. *Int J Health Educ* 1981;24(4):271-9.
215. Mogielnicki RP, Neslin S, Dulac J, Balestra D, Gillie E, Corson J. Tailored media can enhance the success of smoking cessation clinics. *J Behav Med* 1986;9(2):141-61.
216. Platt S, Tannahill A, Watson J, Fraser E. Effectiveness of antismoking telephone helpline: follow up survey. *BMJ* 1997;314(7091):1371-5.
217. Shuster GF, Utz SW, Merwin E. Implementation and outcomes of a community-based self-help smoking cessation program. *J Community Health Nurs* 1996;13(3):187-98.
218. Mudde AN, de Vries H, Strecher VJ. Cost-effectiveness of smoking cessation modalities: comparing apples with oranges? *Prev Med* 1996;25(6):708-16.
219. Ratcliffe J, Cairns J, Platt S. Cost effectiveness of a mass media-led anti-smoking campaign in Scotland. *Tob Control* 1997;6(2):104-10.
220. Fiscella K, Franks P. Cost-effectiveness of the transdermal nicotine patch as an adjunct to physicians' smoking cessation counseling. *JAMA* 1996;275(16):1247-51.
221. Novotny TE, Siegel MB. California's tobacco control saga. *Health Aff* 1996;15(1):58-72.
222. Begay ME, Traynor M, Glantz SA. The tobacco industry, state politics, and tobacco education in California. *Am J Public Health* 1993;83(9):1214-21.
223. Cummings KM, Sciandra R, Markello S. Impact of a newspaper mediated quit smoking program. *Am J Public Health* 1987;77(11):1452-3.
224. Flay BR, Gruder CL, Warnecke RB, Jason LA, Peterson P. One year follow-up of the Chicago televised smoking cessation program. *Am J Public Health* 1989;79(10):1377-80.

225. Jason LA, Gruder CL, Martino S, Flay BR, Warnecke R, Thomas N. Work site group meetings and the effectiveness of a televised smoking cessation intervention. *Am J Community Psychol* 1987;15(1):57-72.

226. Jason LA, Tait E, Goodman D, Buckenberger L, Gruder CL. Effects of a televised smoking cessation intervention among low-income and minority smokers. *Am J Community Psychol* 1988;16(6):863-76.

227. Millar WJ, Naegele BE. Time to quit: community involvement in smoking cessation. *Can J Public Health* 1987;78(2):109-14.

228. Mudde AN, de Vries H. The reach and effectiveness of a national mass media-led smoking cessation campaign in The Netherlands. *Am J Public Health* 1999;89(3):346-50.

229. Salina D, Jason LA, Hedeker D, et al. A follow-up of a media-based, worksite smoking cessation program. *Am J Community Psychol* 1994;22(2):257-71.

230. Sutton S, Hallett R. Randomized trial of brief individual treatment for smoking using nicotine chewing gum in a workplace setting. *Am J Public Health* 1987;77(9):1210-1.

231. Warnecke RB, Langenberg P, Wong SC, Flay BR, Cook TD. The second Chicago televised smoking cessation program: a 24-month follow-up. *Am J Public Health* 1992;82(6):835-40.

232. Ben-Sira Z. The health promoting function of mass media and reference groups: motivating or reinforcing of behavior change. *Soc Sci Med* 1982;16(7):825-34.

233. Danaher BG, Berkanovic E, Gerber B. Mass media based health behavior change: televised smoking cessation program. *Addict Behav* 1984;9(3):245-53.

234. Korhonen HJ, Niemensivu H, Piha T, et al. National TV smoking cessation program and contest in Finland. *Prev Med* 1992;21(1):74-87.

235. Leroux RS, Miller ME. Electronic media-based smoking cessation clinic in the USA. *Hygie* 1983;2(1):23-37.

236. Ryan B, Coffin K, Smillie C, Porter K. Smoking cessation in Nova Scotia: results of the Time to Quit program. *Can J Public Health* 1990;81(2):166-7.

237. Sallis JF, Flora JA, Fortmann SP, Taylor CB, Maccoby N. Mediated smoking cessation programs in the Stanford Five-City Project. *Addict Behav* 1985;10(4):441-3.

238. Thompson B, Curry SJ. Characteristics and predictors of participation and success in a televised smoking cessation activity. *Am J Health Promot* 1994;8(3):175-7.

239. Valois RF, Adams KG, Kammermann SK. One-year evaluation results from Cable-Quit: a community cable television smoking cessation pilot program. *J Behav Med* 1996;19(5):479-99.

240. Wewers ME, Ahijevych K, Page JA. Evaluation of a mass media community smoking cessation campaign. *Addict Behav* 1991;16(5):289-94.

241. Harvey M, Tuffin A, Tuffin K. An evaluation of a community based antismoking campaign. *N Z Med J* 1990;103(890):243-5.

242. Gruder CL, Warnecke RB, Jason LA, Flay BR, Peterson P. A televised, self-help, cigarette smoking cessation intervention. *Addict Behav* 1990;15(6):505-16.

243. Gruder CL, Mermelstein RJ, Kirkendol S, et al. Effects of social support and relapse prevention training as adjuncts to a televised smoking-cessation intervention. *J Consult Clin Psychol* 1993;61(1):113-20.

244. Jason L, Gruder CL, Buckenberger L, et al. A 12-month follow-up of a worksite smoking cessation intervention. *Health Educ Res* 1987;2(3):185-94.

245. Jason LA, Lesowitz T, Michaels M, et al. A worksite smoking cessation intervention involving the media and incentives. *Am J Community Psychol* 1989;17(6):785–99.
246. Kviz FJ, Crittenden KS, Belzer LJ, Warnecke RB. Psychosocial factors and enrollment in a televised smoking cessation program. *Health Educ Q* 1991;18(4):445–61.
247. Kviz FJ, Crittenden KS, Clark MA, Madura KJ, Warnecke RB. Buddy support among older smokers in a smoking cessation program. *J Aging Health* 1994;6(2):229–54.
248. McFall SL, Michener A, Rubin D, et al. The effects and use of maintenance newsletters in a smoking cessation intervention. *Addict Behav* 1993;18(2):151–8.
249. Smillie C, Coffin K, Porter K, Ryan B. Primary health care through a community based smoking-cessation program. *J Community Health* 1988;13(3):156–70.
250. Warnecke RB, Langenberg P, Gruder CL, Flay BR, Jason LA. Factors in smoking cessation among participants in a televised intervention. *Prev Med* 1989;18(6):833–46.
251. Warnecke RB, Flay BR, Kviz FJ, et al. Characteristics of participants in a televised smoking cessation intervention. *Prev Med* 1991;20(3):389–403.
252. Resnicow K, Vaughan R, Futterman R, et al. A self-help smoking cessation program for inner-city African Americans: results from the Harlem Health Connection Project. *Health Educ Behav* 1997;24(2):201–17.
253. Centers for Disease Control and Prevention. Impact of promotion of the Great American Smokeout and availability of over-the-counter nicotine medications, 1996. *MMWR* 1997;46(37):867–71.
254. Chapman S, Smith W, Mowbray G, Hugo C, Egger G. Quit and win smoking cessation contests: how should effectiveness be evaluated? *Prev Med* 1993;22(3):423–32.
255. Chapman S, Smith W. Deception in quit smoking lottery entrants. *Am J Health Promot* 1994;8:328–30.
256. Donovan RJ, Fisher DA, Armstrong BK. “Give it away for a day”: an evaluation of Western Australia’s first smoke free day. *Community Health Stud* 1984;8(3):301–6.
257. Elder JP, McGraw SA, Rodrigues A, et al. Evaluation of two community-wide smoking cessation contests. *Prev Med* 1987;16(2):221–34.
258. Elder JP, Campbell NR, Mielchen SD, Hovell MF, Litrownik AJ. Implementation and evaluation of a community-sponsored smoking cessation contest. *Am J Health Promot* 1991;5(3):200–7.
259. Glasgow RE, Klesges RC, Mizes JS, Pechacek TF. Quitting smoking: strategies used and variables associated with success in a stop-smoking contest. *J Consult Clin Psychol* 1985;53(6):905–12.
260. Gritz ER, Carr CR, Marcus AC. Unaided smoking cessation: Great American Smokeout and New Year’s Day quitters. Special Issue: Clinical research issues in psychosocial oncology. *J Psychosoc Oncol* 1988;6(3–4):217–34.
261. Hantula D, Stillman FA, Waranch HR. Can a mass media campaign modify tobacco smoking in a large organization? Evaluation of the Great American Smokeout in an urban hospital. *J Organ Behav Manage* 1992;13(1):33–47.
262. King AC, Flora JA, Fortmann SP, Taylor CB. Smokers’ challenge: immediate and long-term findings of a community smoking cessation contest. *Am J Public Health* 1987;77(10):1340–1.
263. Korhonen T, Urjanheimo E, Mannonen P, Korhonen HJ, Uutela A, Puska P. Quit and Win campaigns as a long-term anti-smoking intervention in North Karelia and other parts of Finland. *Tob Control* 1999;8:175–81.

264. Lando HA, Loken B, Howard-Pitney B, Pechacek T. Community impact of a localized smoking cessation contest. *Am J Public Health* 1990;80(5):601-3.

265. Lando HA, Pirie PL, Dusich KH, Elsen C, Bernards J. Community incorporation of quit and win contests in Bloomington, Minnesota. *Am J Public Health* 1995;85(2):263-4.

266. Leinweber CE, Macdonald JM, Campbell HS. Community smoking cessation contests: an effective public health strategy. *Can J Public Health* 1994;85(2):95-8.

267. Tillgren P, Haglund BJ, Ainetdin T, Holm LE. Who is a successful quitter? One-year follow-up of a National Tobacco Quit and Win Contest in Sweden. *Scand J Soc Med* 1995;23(3):193-201.

268. Resnicow K, Futterman R, Weston RE, et al. Smoking prevalence in Harlem, New York. *Am J Health Promot* 1996;10(5):343-6.

269. Resnicow K, Royce J, Vaughan R, Orlandi MA, Smith M. Analysis of a multi-component smoking cessation project: what worked and why. *Prev Med* 1997;26(3):373-81.

270. Tillgren P, Haglund BJ, Gilljam H, Holm LE. A tobacco quit and win model in the Stockholm cancer prevention programme. *Eur J Cancer Prev* 1992;1(5):361-6.

271. Chang HC, Zimmerman LH, Beck JM. Impact of chart reminders on smoking cessation practices of pulmonary physicians. *Am J Respir Crit Care Med* 1995;152(3):984-7.

272. Dietrich AJ, O'Conner GT, Keller A, Carney PA, Levy D, Whaley FS. Cancer: improving early detection and prevention. A community practice randomised trial. *BMJ* 1992;304:687-91.

273. Robinson MD, Laurent SL, Little JJ. Including smoking status as a new vital sign: it works. *J Fam Pract* 1995;40(6):556-61.

274. Hahn DL, Berger MG. Implementation of a systematic health maintenance protocol in a private practice. *J Fam Pract* 1990;31(5):492-502.

275. Strecher VJ, O'Malley MS, Villagra VG, et al. Can residents be trained to counsel patients about quitting smoking? Results from a randomized trial. *J Gen Intern Med* 1991;6(1):9-17.

276. Rosser WW, McDowell I, Newell C. Use of reminders for preventive procedures in family medicine. *Can Med Assoc J* 1991;145(7):807-13.

277. Spencer E, Swanson T, Hueston WJ, Edberg DL. Tools to improve documentation of smoking status. Continuous quality improvement and electronic medical records. *Arch Fam Med* 1999;8(1):18-22.

278. Fiore MC, Jorenby DE, Schensky AE, Smith SS, Bauer RR, Baker TB. Smoking status as the new vital sign: effect on assessment and intervention in patients who smoke. *Mayo Clin Proc* 1995;70(3):209-13.

279. Cohen SJ, Stookey GK, Katz BP, Drook CA, Smith DM. Encouraging primary care physicians to help smokers quit. A randomized, controlled trial. *Ann Intern Med* 1989;110(8):648-52.

280. Cohen SJ, Stookey GK, Katz BP, Drook CA, Christen AG. Helping smokers quit: a randomized controlled trial with private practice dentists. *J Am Dent Assoc* 1989;118:41-5.

281. Conger B, Nelson EC, Dietrich AJ, et al. Effectiveness of physician antismoking advice. *Am J Prev Med* 1987;3(4):223-6.

282. Cummings SR, Richard RJ, Duncan CL, et al. Training physicians about smoking cessation: a controlled trial in private practice. *J Gen Intern Med* 1989;4(6):482–9.
283. Cummings SR, Coates TJ, Richard RJ, et al. Training physicians in counseling about smoking cessation. A randomized trial of the “Quit for Life” program. *Ann Intern Med* 1989;110(8):640–7.
284. Davies BL, Matte-Lewis L, O’Connor AM, Dulberg CS, Drake ER. Evaluation of the “Time to Quit” self-help smoking cessation program. *Can J Public Health* 1992; 83(1):19–23.
285. Duncan C, Stein MJ, Cummings SR. Staff involvement and special follow-up time increase physicians’ counseling about smoking cessation: a controlled trial. *Am J Public Health* 1991;81(7):899–901.
286. Gilbert JR, Wilson DM, Singer J, et al. A family physician smoking cessation program: an evaluation of the role of follow-up visits. *Am J Prev Med* 1992;8(2):91–5.
287. Goldberg DN, Hoffman AM, Farinha MF, et al. Physician delivery of smoking-cessation advice based on the stages-of-change model. *Am J Prev Med* 1994;10(5): 267–74.
288. Hartmann KE, Thorp MJ, Pahel-Short L, Koch MA. A randomized controlled trial of smoking cessation intervention in pregnancy in an academic clinic. *Obstet Gynecol* 1996;87(4):621–6.
289. Hollis JF, Lichtenstein E, Vogt TM, Stevens VJ, Biglan A. Nurse-assisted counseling for smokers in primary care. *Ann Intern Med* 1993;118:521–5.
290. Hovell MF, Slymen DJ, Jones JA, et al. An adolescent tobacco-use prevention trial in orthodontic offices. *Am J Public Health* 1996;86(12):1760–6.
291. Kottke TE, Brekke ML, Solberg LI, Hughes JR. A randomized trial to increase smoking intervention by physicians: Doctors Helping Smokers, Round I. *JAMA* 1989; 261(14):2101–6.
292. Kottke TE, Solberg LI, Brekke ML, Conn SA, Maxwell P, Brekke MJ. A controlled trial to integrate smoking cessation advice into primary care practice: Doctors Helping Smokers, Round III. *J Fam Pract* 1992;34(6):701–8.
293. Manfredi C, Crittenden KS, Warnecke RB, Engler J, Cho YI, Shaligram C. Evaluation of a motivational smoking cessation intervention for women in public health clinics. *Prev Med* 1999;28:51–60.
294. McIlvain HE, Susman JL, Manners MA, Davis CM, Gilbert CS. Improving smoking cessation counseling by family practice residents. *J Fam Pract* 1992;34(6):745–9.
295. McPhee S, Bird J, Fordham D, Rodnick J. Promoting cancer prevention activities by primary care physicians: results of a randomized, controlled trial. *JAMA* 1991;266: 538–44.
296. Morgan GD, Noll EL, Orleans CT, Rimer BK, Amfoh K, Bonney G. Reaching midlife and older smokers: tailored interventions for routine medical care. *Prev Med* 1996;25(3):346–54.
297. Ockene JK, Kristeller J, Pbert L, et al. The physician-delivered smoking intervention project: can short-term interventions produce long-term effects for a general outpatient population? *Health Psychol* 1994;13(3):278–81.
298. Petersen L, Handel J, Kotch J, Podedworny T, Rosen A. Smoking reduction during pregnancy by a program of self-help and clinical support. *Obstet Gynecol* 1992; 79(6):924–30.

299. Richmond R, Mendelsohn C, Kehoe L. Family physicians' utilization of a brief smoking cessation program following reinforcement contact after training: a randomized trial. *Prev Med* 1998;27(1):77-83.

300. Rigotti NA, Arnsten JH, McKool KM, Wood-Reid KM, Pasternak RC, Singer DE. Efficacy of a smoking cessation program for hospital patients. *Arch Intern Med* 1997;157(22):2653-60.

301. Rodney WM, Nutter D, Widoff B. Recording patients' consumption of social drugs in a family medicine residency: a longitudinal study. *Fam Pract* 1985;2(2):86-90.

302. Secker-Walker RH, Solomon LJ, Flynn BS, Skelly JM, Mead PB. Reducing smoking during pregnancy and postpartum: physician's advice supported by individual counseling. *Prev Med* 1998;27:422-30.

303. Sinclair HK, Bond CM, Lennox AS, Silcock J, Windfield AJ, Donnan PT. Training pharmacists and pharmacy assistants in the stage-of-change model of smoking cessation: a randomised controlled trial in Scotland. *Tob Control* 1998;7:253-61.

304. Stevens VJ, Severson H, Lichtenstein E, Little SJ, Leben J. Making the most of a teachable moment: a smokeless-tobacco cessation intervention in the dental office. *Am J Public Health* 1995;85(2):231-5.

305. Wall MA, Severson HH, Andrews JA, Lichtenstein E, Zoref L. Pediatric office-based smoking intervention: impact on maternal smoking and relapse. *Pediatrics* 1995;96(4 pt 1):622-8.

306. Wilson DM, Taylor DW, Gilbert JR, et al. A randomized trial of a family physician intervention for smoking cessation. *JAMA* 1988;260(11):1570-4.

307. Windsor RA, Lowe JB, Perkins LL, et al. Health education for pregnant smokers: its behavioral impact and cost benefit. *Am J Public Health* 1993;83(2):201-6.

308. Cohen SJ, Christen AG, Katz BP, et al. Counseling medical and dental patients about cigarette smoking: the impact of nicotine gum and chart reminders. *Am J Public Health* 1987;77(3):313-6.

309. Hebert JR, Kristeller J, Ockene JK, et al. Patient characteristics and the effect of three physician-delivered smoking interventions. *Prev Med* 1992;21(5):557-73.

310. Hollis JF, Lichtenstein E, Mount K, Vogt TM, Stevens VJ. Nurse-assisted smoking counseling in medical settings: minimizing demands on physicians. *Prev Med* 1991;20(4):497-507.

311. Kottke TE, Solberg LI, Conn S, et al. A comparison of two methods to recruit physicians to deliver smoking cessation interventions. *Arch Intern Med* 1990;150(7):1477-81.

312. Lichtenstein E, Hollis J. Patient referral to a smoking cessation program: who follows through? *J Fam Pract* 1992;34(6):739-44.

313. Lindsay-McIntyre E, Wilson D, Best JA, et al. The impact of a continuing education package for smoking cessation on physicians' clinical behavior and patient smoking. *Proc Annu Conf Res Med Educ* 1987;26:14-9.

314. Lindsay EA, Wilson D, Best JA, Singer J, Gilbert JR, Taylor DW. A randomized trial of physician training for smoking cessation. *Am J Health Promot* 1989;3(3):11-8.

315. Little SJ, Stevens VJ, Severson HH, Lichtenstein E. Effective smokeless tobacco intervention for dental hygiene patients. *J Dent Hyg* 1992;66(4):185-90.

316. Manfredi C, Crittenden KS, Cho YI, Engler J, Warnecke RB. Minimal smoking cessation interventions in prenatal, family planning, and well-child public health clinics. *Am J Public Health* 2000;90:423-7.

317. Manfredi C, Crittenden KS, Cho YI, Engler J, Warnecke RB. The effect of a structured smoking cessation program, independent of exposure to existing interventions. *Am J Public Health* 2000;90:751–6.

318. Ockene JK, Kristeller J, Goldberg R, et al. Increasing the efficacy of physician-delivered smoking interventions: a randomized clinical trial. *J Gen Intern Med* 1991; 6(1):1–8.

319. Rosal MC, Ockene JK, Hurley TG, Kalan K, Hebert JR. Effectiveness of nicotine-containing gum in the Physician-Delivered Smoking Intervention Study. *Prev Med* 1998; 27(2):262–7.

320. Secker-Walker RH, Solomon LJ, Flynn BS, et al. Training obstetric and family practice residents to give smoking cessation advice during prenatal care. *Am J Obstet Gynecol* 1992;166(5):1356–63.

321. Whitlock EP, Vogt TM, Hollis JF, Lichtenstein E. Does gender affect response to a brief clinic-based smoking intervention? *Am J Med* 1997;13(3):159–66.

322. Windsor RA, Lowe JB, Artz L, Contreras L. Smoking cessation and pregnancy intervention trial: preliminary mid-trial results. *Prog Clin Biol Res* 1990;339:107–17.

323. Allen B, Pederson LL, Leonard EH. Effectiveness of physicians-in-training counseling for smoking cessation in African Americans. *J Natl Med Assoc* 1998;90(10): 597–604.

324. Kosower E, Ernst A, Taub B, Berman N, Andrews J, Seidel J. Tobacco prevention education in a pediatric residency program. *Arch Pediatr Adolesc Med* 1995;149(4): 430–5.

325. Leshan LA, Fitzsimmons M, Marbella A, Gottlieb M. Increasing clinical prevention efforts in a family practice residency program through CQI methods. *J Qual Improv* 1997;23(7):391–400.

326. Narce-Valente S, Kligman EW. Increasing physician screening and counseling for passive smoking. *J Fam Pract* 1992;34(6):722–8.

327. Russell MA, Stapleton JA, Jackson PH, Hajek P, Belcher M. District programme to reduce smoking: effect of clinic supported brief intervention by general practitioners. *BMJ (Clinical Research Ed)* 1987;295:1240–4.

328. Solberg LI, Maxwell PL, Kottke TE, Gepner GJ, Brekke ML. A systematic primary care office-based smoking cessation program. *J Fam Pract* 1990;30(6):647–54.

329. Allen SS, Bland CJ, Dawson SJ. A mini-workshop to train medical students to use a patient-centered approach to smoking cessation. *Am J Prev Med* 1990;6(1): 28–33.

330. Carney PA, Dietrich AJ, Freeman DJ, Mott LA. A standardized-patient assessment of a continuing medical education program to improve physicians' cancer-control clinical skills. *Acad Med* 1995;70(1):52–8.

331. Cockburn J, Ruth D, Silagy C, et al. Randomised trial of three approaches for marketing smoking cessation programmes to Australian general practitioners. *BMJ* 1992; 304(6828):691–4.

332. Giovino GA, Cummings KM, Koenigsberg MR, Sciandra RC. An evaluation of a physician training program on patient smoking cessation. *Prog Clin Biol Res* 1990;339: 27–48.

333. Klein JD, Portilla M, Goldstein A, Leininger L. Training pediatric residents to prevent tobacco use. *Pediatrics* 1995;96(2 pt 1):326–30.

334. Ockene JK, Lindsay EA, Hymowitz N, et al. Tobacco control activities of primary-

care physicians in the Community Intervention Trial for Smoking Cessation. COMMIT Research Group. *Tob Control* 1997;6(suppl 2):S49-S56.

335. Quirk M, Ockene J, Kristeller J, et al. Training family practice and internal medicine residents to counsel patients who smoke: improvement and retention of counseling skills. *Fam Med* 1991;23(2):108-11.

336. Roche AM, Eccleston P, Sanson-Fisher R. Teaching smoking cessation skills to senior medical students: a block-randomized controlled trial of four different approaches. *Prev Med* 1996;25(3):251-8.

337. Royce JM, Gorin SS, Edelman B, Rendino-Perrone R, Orlandi MA. Student nurses and smoking cessation. *Prog Clin Biol Res* 1990;339:49-71.

338. Ward J, Sanson-Fisher R. Does a 3-day workshop for family medicine trainees improve preventive care? A randomized control trial. *Prev Med* 1996;25(6):741-7.

339. Dietrich AJ, Barrett J, Levy D, Carney-Gersten P. Impact of an educational program on physician cancer control knowledge and activities. *Am J Prev Med* 1990;6(6):346-52.

340. Ockene JK, Quirk ME, Goldberg RJ, et al. A residents' training program for the development of smoking intervention skills. *Arch Intern Med* 1988;148(5):1039-45.

341. Anonymous. The impact of providing physicians with quit-smoking materials for smoking patients. *CA Cancer J Clin* 1981;31(2):75-8.

342. Chu F, Day R. Smoking recognition by family physicians. *J Fam Pract* 1981;12(4):657-60.

343. Dunkley J. Training midwives to help pregnant women stop smoking. *Nurs Times* 1997;93(5):64-6.

344. Hall MN, Pettice YJ, Robinson MD, Alexander M. Teaching smoking cessation to family practice residents: an experiential approach. *Fam Med* 1996;28(5):331-6.

345. Leininger LS, Earp JA. The effect of training staff in office-based smoking cessation counseling. *Patient Educ Couns* 1993;20(1):17-25.

346. Montner P, Bennett G, Brown C. An evaluation of a smoking cessation training program for medical residents in an inner-city hospital. *J Natl Med Assoc* 1994;86(9):671-5.

347. Patterson J, Fried RA, Nagle JP. Impact of a comprehensive health promotion curriculum on physician behavior and attitudes. *Am J Prev Med* 1989;5(1):44-9.

348. Wood GJ, Cecchini JJ, Nathason N, Hiroshige K. Office-based training in tobacco cessation for dental professionals. *J Am Dent Assoc* 1997;128(2):216-24.

349. Bonevski B, Sanson-Fisher RW, Campbell E, Carruthers A, Reid AL, Ireland M. Randomized controlled trial of a computer strategy to increase general practitioner preventive care. *Prev Med* 1999;29(6 pt 1):478-86.

350. Kern D, Harris W. Use of an outpatient medical record audit to achieve educational objectives. *J Gen Intern Med* 1990;5:218-24.

351. Shank JC, Powell T, Llewelyn J. A five-year demonstration project association with improvement in physician health maintenance behavior. *Fam Med* 1989;21(4):273-8.

352. Ewart CK, Li VC, Coates TJ. Increasing physicians' antismoking influence by applying an inexpensive feedback technique. *J Med Educ* 1983;58(6):468-73.

353. Frame P, Kowulich B, Llewellyn A. Improving physician compliance with a health maintenance protocol. *J Fam Pract* 1984;19:341-4.

354. Cox JL, McKenna JP. Nicotine gum: does providing it free in a smoking cessation program alter success rates? *J Fam Pract* 1990;31(3):278-80.

355. Curry SJ, Grothaus LC, McAfee T, Pabiniak C. Use and cost effectiveness of smoking-cessation services under four insurance plans in a health maintenance organization. *N Engl J Med* 1998;339(10):673–9.
356. Hughes JR, Wadland WC, Fenwick JW, Lewis J, Bickel WK. Effect of cost on the self-administration and efficacy of nicotine gum: a preliminary study. *Prev Med* 1991;20(4):486–96.
357. Johnson RE, Hollis JF, Stevens VJ, Woodson GT. Patterns of nicotine gum use in a health maintenance organization. *DICP* 1991;25:730–5.
358. Schauflier HH, McMenemy S, Olson K, Boyce-Smith G, Rideout JA, Kamil J. Variations in treatment benefits influence smoking cessation: results of a randomized controlled trial. *Tob Control* 2001;10(2):175–80.
359. Cromwell J, Bartosch WJ, Fiore MC, Hasselblad V, Baker T. Cost-effectiveness of the clinical practice recommendations in the AHCPR guideline for smoking cessation. Agency for Health Care Policy and Research. *JAMA* 1997;278(21):1759–66.
360. Sofian NS, McAfee T, Wilson J, Levan S. Telephone smoking cessation intervention: the Free and Clear Program. *HMO Pract* 1995;9(3):144–6.
361. Curry SJ, McBride C, Grothaus LC, Louie D, Wagner EH. A randomized trial of self-help materials, personalized feedback, and telephone counseling with nonvolunteer smokers. *J Consult Clin Psychol* 1995;63(6):1005–14.
362. Decker BD, Evans RG. Efficacy of a minimal contact version of a multimodal smoking cessation program. *Addict Behav* 1989;14(5):487–91.
363. Gebauer C, Kwo CY, Haynes EF, Wewers ME. A nurse-managed smoking cessation intervention during pregnancy. *J Obstet Gynecol Neonatal Nurs* 1998;27(1):47–53.
364. Glasgow RE, Whitlock EP, Eakin EG, Lichtenstein E. A brief smoking cessation intervention for women in low-income planned parenthood clinics. *Am J Public Health* 2000;90(5):786–9.
365. Johnson JL, Budz B, Mackay M, Miller C. Evaluation of a nurse-delivered smoking cessation intervention for hospitalized patients with cardiac disease. *Heart Lung* 1999;28(1):55–64.
366. Lando HA, Hellerstedt WL, Pirie PL, McGovern PG. Brief supportive telephone outreach as a recruitment and intervention strategy for smoking cessation. *Am J Public Health* 1992;82(1):41–6.
367. Lando HA, Pirie PL, Roski J, McGovern PG, Schmid LA. Promoting abstinence among relapsed chronic smokers: the effect of telephone support. *Am J Public Health* 1996;86(12):1786–90.
368. Lando HA, Rolnick S, Klevan D, Roski J, Cherney L, Lauger G. Telephone support as an adjunct to transdermal nicotine in smoking cessation. *Am J Public Health* 1997;87(10):1670–4.
369. Lewis SF, Piasecki TM, Fiore MC, Anderson JE, Baker TB. Transdermal nicotine replacement for hospitalized patients: a randomized clinical trial. *Prev Med* 1998;27(2):296–303.
370. McBride CM, Curry SJ, Lando HA, Pirie PL, Grothaus LC, Nelson JC. Prevention of relapse in women who quit smoking during pregnancy. *Am J Public Health* 1999;89(5):706–11.
371. McBride CM, Scholes D, Grothaus LC, Curry SJ, Ludman E, Albright J. Evaluation of a minimal self-help smoking cessation intervention following cervical cancer screening. *Prev Med* 1999;29(2):133–8.

372. Miller N, Smith P, DeBusk RF, Sobel DS, Taylor CB. Smoking cessation in hospitalized patients: results of a randomized trial. *Arch Fam Med* 1997;157:409–15.

373. Orleans CT, Schoenbach VJ, Wagner EH, et al. Self-help quit smoking interventions: effects of self-help materials, social support instructions, and telephone counseling. *J Consult Clin Psychol* 1991;59(3):439–48.

374. Orleans CT, Boyd NR, Bingle R, et al. A self-help intervention for African American smokers: tailoring cancer information service counseling for a special population. *Prev Med* 1998;27(5 (pt 2)):S61–S70.

375. Ossip-Klein DJ, Giovino GA, Megahed N, et al. Effects of a smoker's hotline: results of a 10-county self-help trial. *J Consult Clin Psychol* 1991;59(2):325–32.

376. Ossip-Klein DJ, Carosella AM, Krusch DA. Self-help interventions for older smokers. *Tob Control* 1997;6(3):188–93.

377. Prochaska JO, DiClemente CC, Velicer WF, Rossi JS. Standardized, individualized, interactive, and personalized self-help programs for smoking cessation. *Health Psychol* 1993;12(5):399–405.

378. Sexton M, Hebel JR. A clinical trial of change in maternal smoking and its effect on birth weight. *JAMA* 1984;251(7):911–5.

379. Simon JA, Solkowitz SN, Carmody TP, Browner WS. Smoking cessation after surgery, a randomized trial. *Arch Fam Med* 1997;157:1371–6.

380. Stevens VJ, Glasgow RE, Hollis JF, Lichtenstein E, Vogt TM. A smoking-cessation intervention for hospital patients. *Med Care* 1993;31(1):65–72.

381. Thompson B, Kinne S, Lewis FM, Woolridge JA. Randomized telephone smoking-intervention trial initially directed at blue-collar workers. *J Natl Cancer Inst Monogr* 1993;14:105–12.

382. Weissfeld JL, Holloway JL. Treatment for cigarette smoking in a Department of Veterans Affairs outpatient clinic. *Arch Intern Med* 1991;151(5):973–7.

383. Westman EC, Levin ED, Rose JE. The nicotine patch in smoking cessation. A randomized trial with telephone counseling. *Arch Intern Med* 1993;153(16):1917–23.

384. Zhu SH, Stretch V, Balabanis M, Rosbrook B, Sadler G, Pierce JP. Telephone counseling for smoking cessation: effects of single-session and multiple-session interventions. *J Consult Clin Psychol* 1996;64(1):202–11.

385. Britt J, Curry SJ, McBride C, Grothaus L, Louie D. Implementation and acceptance of outreach telephone counseling for smoking cessation with nonvolunteer smokers. *Health Educ Q* 1994;21(1):55–68.

386. Glasgow RE, Lando H, Hollis J, McRae SG, La Chance PA. A stop-smoking telephone help line that nobody called. *Am J Public Health* 1993;83(2):252–3.

387. Hebel JR, Nowicki P, Sexton M. The effect of antismoking intervention during pregnancy: an assessment of interactions with maternal characteristics. *Am J Epidemiol* 1985;122(1):135–48.

388. Kinne S, Thompson B, Wooldridge JA. Response to a telephone smoking information line. *Am J Health Promot* 1991;5(6):410–3.

389. Meenan RT, Stevens VJ, Hornbrook MC, et al. Cost-effectiveness of a hospital-based smoking cessation intervention. *Med Care* 1998;36(5):670–8.

390. Taylor CB, Miller N, Herman S, et al. A nurse-managed smoking cessation program for hospitalized smokers. *Am J Public Health* 1996;86(11):1557–60.

391. Vogt TM, Lichtenstein E, Ary D, et al. Integrating tobacco intervention into a

health maintenance organization: the TRACC program. Special Issue: Smoking. *Health Educ Res* 1989;4(1):125–35.

392. Amos A, White DA, Elton RA. Is a telephone helpline of value to the workplace smoker? *Occup Med* 1995;45(5):234–8.

393. Bachman DS. Smoking cessation via telephone counseling and tools for helping your patients quit smoking once and for all. *J Ark Med Soc* 1997;94(4):147–8.

394. Dubren R. Self-reinforcement by recorded telephone messages to maintain non-smoking behavior. *J Consult Clin Psychol* 1977;45(3):358–60.

395. Glasgow RE, Hollis JF, McRae SG, Lando HA, LaChance P. Providing an integrated program of low intensity tobacco cessation services in a health maintenance organization. *Health Educ Res* 1991;6(1):87–9.

396. Racelis MC, Lombardo K, Verdin J. Impact of telephone reinforcement of risk reduction education on patient compliance. *J Vasc Nurs* 1998;16(1):16–20.

397. Croghan IT, Offord KP, Evans RW, et al. Cost-effectiveness of treating nicotine dependence: the Mayo Clinic experience. *Mayo Clin Proc* 1997;72(10):917–24.

398. Ershoff DH, Aaronson NK, Danaher BG, Wasserman FW. Behavioral, health, and cost outcomes of an HMO-based prenatal health education program. *Public Health Rep* 1983;98(6):536–47.

399. Krumholz HM, Cohen BJ, Tsevat J, Pasternak RC, Weinstein MC. Cost-effectiveness of a smoking cessation program after myocardial infarction. *J Am Coll Cardiol* 1993;22(6):1697–702.

400. Marks JS, Koplan JP, Hogue CJ, Dalmat ME. A cost-benefit/cost-effectiveness analysis of smoking cessation for pregnant women. *Am J Prev Med* 1990;6(5):282–9.

401. U.S. Office on Smoking and Health. The health consequences of involuntary smoking. A report of the Surgeon General. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, Center for Health Promotion and Education, Office on Smoking and Health, 1986. DHHS Pub No (CDC) 87-8398.

402. Burns DM, Axelrad R, Bal D, et al. Report of the Tobacco Policy Research Study Group on Smoke-Free Indoor Air Policies. *Tob Control* 1992;1(suppl):S14–S18.

403. Fielding JE. Banning worksite smoking. *Am J Public Health* 1986;76(8):957–9.

404. Pierce JP, Naquin M, Gilpin E, Giovino G, Mills S, Marcus S. Smoking initiation in the United States: a role for worksite and college smoking bans. *J Natl Cancer Inst* 1991;83(14):1009–13.

405. Becker DM, Conner HF, Waranch HR, et al. The impact of a total ban on smoking in the Johns Hopkins Children's Center. *JAMA* 1989;262(6):799–802.

406. Borland R, Pierce JP, Burns DM, Gilpin E, Johnson M, Bal D. Protection from environmental tobacco smoke in California: the case for a smoke-free workplace. *JAMA* 1992;268(6):749–52.

407. Broder I, Pilger C, Corey P. Environment and well-being before and following smoking ban in office building. *Can J Public Health* 1993;84(4):254–8.

408. Brownson RC, Davis J, Jackson-Thompson J, Wilkerson J. Environmental tobacco smoke awareness and exposure: impact of a statewide clean indoor air law and the report of the U.S. Environmental Protection Agency. *Tob Control* 1995;4:132–8.

409. Etter J, Ronchi A, Perneger TV. Short-term impact of a university based smoke free campaign. *J Epidemiol Community Health* 1999;53:710–5.

410. Gottlieb NH, Eriksen MP, Lovato CY, Weinstein RP, Green LW. Impact of a restrictive work site smoking policy on smoking behavior, attitudes, and norms. *J Occup Med* 1990;32(1):16–23.

411. Millar WJ. Evaluation of the impact of smoking restrictions in a government work setting. *Can J Public Health* 1988;79(5):379–82.

412. Mullooly JP, Schuman KL, Stevens VJ, Glasgow RE, Vogt TM. Smoking behavior and attitudes of employees of a large HMO before and after a work site ban on cigarette smoking. *Public Health Rep* 1990;105(6):623–8.

413. Patten CA, Pierce JP, Cavin SW, Berry C, Kaplan R. Progress in protecting non-smokers from environmental tobacco smoke in California workplaces. *Tob Control* 1995;4:139–44.

414. Stillman FA, Becker DM, Swank RT, et al. Ending smoking at the Johns Hopkins Medical Institutions. An evaluation of smoking prevalence and indoor air pollution. *JAMA* 1990;264(12):1565–9.

415. Abernathy T, O'Grady B, Dukeshire S. Changes in ETS following anti-smoking legislation. *Can J Public Health* 1998;89(1):33–4.

416. Batlle E, Boixet M, Agudo A, Almirall J, Salvador T. Tobacco prevention in hospitals: long-term follow-up of a smoking control programme. *Brit J Addict* 1991;86(6):709–17.

417. Borland R, Owen N, Hocking B. Changes in smoking behaviour after a total workplace smoking ban. *Aust J Public Health* 1991;15(2):130–4.

418. Centers for Disease Control. Evaluation of an employee smoking policy—Pueblo, Colorado, 1989–90. *MMWR* 1990;39(38):673–6.

419. Eisner MD, Smith AK, Blanc PD. Bartenders' respiratory health after establishment of smoke-free bars and taverns. *JAMA* 1998;280(22):1909–14.

420. Hammond SK, Sorensen G, Youngstrom R, Ockene JK. Occupational exposure to environmental tobacco smoke. *JAMA* 1995;274(12):956–60.

421. Marcus BH, Emmons K, Abrams DB, et al. Restrictive workplace smoking policies: impact on nonsmokers' tobacco exposure. *J Public Health Policy* 1992;13:42–51.

422. Biener L, Abrams DB, Follick MJ, Dean L. A comparative evaluation of a restrictive smoking policy in a general hospital. *Am J Public Health* 1989;79(2):192–5.

423. Brigham J, Gross J, Stitzer ML, Felch LJ. Effects of a restricted work-site smoking policy on employees who smoke. *Am J Public Health* 1994;84(5):773–8.

424. Jeffery RW, Kelder SH, Forster JL, French SA, Lando HA, Baxter JE. Restrictive smoking policies in the workplace: effects on smoking prevalence and cigarette consumption. *Prev Med* 1994;23(1):78–82.

425. Longo DR, Brownson RC, Johnson JC, et al. Hospital smoking bans and employee smoking behavior: results of a national survey. *JAMA* 1996;275(16):1252–7.

426. Patten CA, Gilpin E, Cavin SW, Pierce JP. Workplace smoking policy and changes in smoking behavior in California: a suggested association. *Tob Control* 1995;4:36–41.

427. Stave GM, Jackson GW. Effect of a total work-site smoking ban on employee smoking and attitudes. *J Occup Med* 1991;33(8):884–90.

428. Bartosch W, Pope G. The economic effect of smoke-free restaurant policies on restaurant business in Massachusetts. *J Public Health Manag Pract* 1999;5:53–62.

429. Centers for Disease Control and Prevention. Assessment of the impact of a 100% smoke-free ordinance on restaurant sales—West Lake Hills, Texas, 1992–1994. *MMWR* 1995;44(19):370–2.

430. Glantz S, Smith L. The effect of ordinances requiring smoke-free restaurants and bars on revenues: a follow-up. *Am J Public Health* 1997;87(10):1687–93.
431. Glantz SA, Smith LR. The effect of ordinances requiring smoke-free restaurants on restaurant sales. *Am J Public Health* 1994;84(7):1081–5.
432. Glantz SA, Charlesworth A. Tourism and hotel revenues before and after passage of smoke-free restaurant ordinances. *JAMA* 1999;281(20):1911–8.
433. Hyland A, Cummings KM, Nauenberg E. Analysis of taxable sales receipts: was New York City's smoke-free air bad for restaurant business? *J Public Health Manag Pract* 1999;5:14–21.
434. Sciacca J, Eckrem M. Effects of a city ordinance regulating smoking in restaurants and retail stores. *J Community Health* 1993;18(3):175–82.
435. Mudarri DH. The costs and benefits of smoking restrictions: an assessment of the Smoke-Free Environment Act of 1993 (H.R. 3434). Government Document 1994; H.R. 3434, April 1994.
436. Ashley MJ, Ferrence R. Reducing children's exposure to environmental tobacco smoke in homes: issues and strategies. *Tob Control* 1998;7:61–5.
437. Greenberg RA, Strecher VJ, Bauman KE, et al. Evaluation of a home-based intervention program to reduce infant passive smoking and lower respiratory illness. *J Behav Med* 1994;17(3):273–90.
438. Farkas AJ, Gilpin E, Distefan JM, Pierce JP. The effects of household and workplace smoking restrictions on quitting behaviors. *Tob Control* 1999;8:261–5.
439. Borland R, Mullins R, Trotter L, White V. Trends in environmental tobacco smoke restrictions in the home in Victoria, Australia. *Tob Control* 1999;8:266–71.
440. Cochrane Collaboration. The Cochrane Library, Issue 4. Available at: www.cochrane.org. Accessed January 6, 2003.
441. National Cancer Policy Board—Institute of Medicine and Commission on Life Sciences—National Research Council. Taking action to reduce tobacco use. Washington, DC: National Academy Press, 1998.