

Alcohol Electronic Screening and Brief Intervention



A Community Guide Systematic Review

Kristin A. Tansil, MSW,¹ Marissa B. Esser, PhD, MPH,¹ Paramjit Sandhu, MD, MPH,¹ Jeffrey A. Reynolds, MPH,¹ Randy W. Elder, PhD, MEd,¹ Rebecca S. Williamson, MPH,¹ Sajal K. Chattopadhyay, PhD,¹ Michele K. Bohm, MPH,¹ Robert D. Brewer, MD, MSPH,² Lela R. McKnight-Eily, PhD,³ Daniel W. Hungerford, DrPH,³ Traci L. Toomey, PhD,⁴ Ralph W. Hingson, ScD, MPH,⁵ Jonathan E. Fielding, MD, MPH, MBA,⁶ and the Community Preventive Services Task Force

Context: Excessive drinking is responsible for one in ten deaths among working-age adults in the U.S. annually. Alcohol screening and brief intervention is an effective but underutilized intervention for reducing excessive drinking among adults. Electronic screening and brief intervention (e-SBI) uses electronic devices to deliver key elements of alcohol screening and brief intervention, with the potential to expand population reach.

Evidence acquisition: Using Community Guide methods, a systematic review of the scientific literature on the effectiveness of e-SBI for reducing excessive alcohol consumption and related harms was conducted. The search covered studies published from 1967 to October 2011. A total of 31 studies with 36 study arms met quality criteria and were included in the review. Analyses were conducted in 2012.

Evidence synthesis: Twenty-four studies (28 study arms) provided results for excessive drinkers only and seven studies (eight study arms) reported results for all drinkers. Nearly all studies found that e-SBI reduced excessive alcohol consumption and related harms: nine study arms reported a median 23.9% reduction in binge-drinking intensity (maximum drinks/binge episode) and nine study arms reported a median 16.5% reduction in binge-drinking frequency. Reductions in drinking measures were sustained for up to 12 months.

Conclusions: According to Community Guide rules of evidence, e-SBI is an effective method for reducing excessive alcohol consumption and related harms among intervention participants. Implementation of e-SBI could complement population-level strategies previously recommended by the Community Preventive Services Task Force for reducing excessive drinking (e.g., increasing alcohol taxes and regulating alcohol outlet density).

(Am J Prev Med 2016;51(5):801–811) Published by Elsevier Inc. on behalf of American Journal of Preventive Medicine

From the ¹Community Guide Branch, Division of Public Health Information Dissemination, Center for Surveillance, Epidemiology, and Laboratory Services, Centers for Disease Control and Prevention (CDC), Atlanta, Georgia;

²Division of Population Health, National Center for Chronic Disease Prevention and Health Promotion, CDC, Atlanta, Georgia; ³National Center on Birth Defects and Developmental Disabilities, CDC, Atlanta, Georgia;

⁴University of Minnesota School of Public Health, Minneapolis, Minnesota;

⁵National Institute of Alcohol Abuse and Alcoholism, Bethesda, Maryland; and ⁶Los Angeles County Department of Public Health, Los Angeles, California

Names and affiliations of Community Preventive Services Task Force members can be found at: www.thecommunityguide.org/about/task-force-members.html.

Address correspondence to: Randy W. Elder, PhD, MEd, Community Guide Branch, Division of Public Health Information Dissemination, Center for Surveillance, Epidemiology, and Laboratory Services, Centers for Disease Control and Prevention, 1600 Clifton Road, MS E-69, Atlanta GA 30329. E-mail: rfe3@cdc.gov.

0749-3797/\$36.00

<http://dx.doi.org/10.1016/j.amepre.2016.04.013>

Context

Excessive alcohol consumption is responsible for approximately 88,000 deaths, including one in ten deaths among working-age adults, in the U.S. each year, costing nearly \$250 billion in 2010.^{1–5} Binge drinking (i.e., four or more drinks for women and five or more drinks for men, per occasion) is responsible for more than half of the deaths, two thirds of the years of potential life lost, and three quarters of the economic costs of excessive drinking.^{5,6} Furthermore, about one in six U.S. adults report binge drinking an average of four times a month, consuming an average of eight drinks per binge episode.⁵ Excessive drinking is also associated with a wide range of health and social problems, including motor vehicle crashes, violence, sexually transmitted diseases, fetal alcohol spectrum disorder, cancers, heart disease, and unintended pregnancy.^{6–9} Yet 90% of adult excessive drinkers do not meet the DSM-IV criteria for alcohol dependence.¹⁰

Alcohol screening and brief intervention (ASBI) is a highly effective strategy for reducing excessive drinking among adults. Traditional ASBI is usually provided in clinical settings, involving screening for excessive alcohol use and providing excessive drinkers with a brief intervention usually lasting 5–15 minutes.^{11–13} Screening tools to assess drinking patterns range from a single-question screen for binge drinking to longer screening tools, such as the Alcohol Use Disorders Identification Test (AUDIT).^{14–16} Those who are found to have alcohol use disorders can be referred to specialized treatment.

In 2004 and 2013, the U.S. Preventive Services Task Force recommended the use of ASBI for adults in primary care settings.¹⁷ In 2006, the National Commission on Prevention Priorities ranked ASBI fourth in a priority listing of clinical preventive services because ASBI was shown to effectively reduce the high preventable burden of excessive drinking and to be cost effective.¹⁸ Since 2010, the Affordable Care Act has required new health insurance plans to cover ASBI without a copayment based on strong scientific evidence of intervention effectiveness.¹⁹ Reducing the proportion of adults who drink excessively and the proportion of adults who engage in binge drinking, and increasing the number of Level I and Level II trauma centers and primary care settings that implement ASBI, are also health objectives in *Healthy People 2020*.²⁰

Despite the scientific evidence of ASBI effectiveness for reducing excessive drinking among adults, it is underutilized in clinical practice.^{21,22} A recent study reported that only 16% of U.S. adults and 25% of adult binge drinkers reported ever discussing alcohol use with a health professional, showing little change since 1997.¹³

Furthermore, only about one third of adults who reported binge drinking ten or more times in the past month had ever discussed their alcohol consumption with a health professional.^{13,23} Common barriers to implementing ASBI in clinical settings include time constraints, lack of training, and self-efficacy.^{13,24}

Alternative methods for delivering key elements of ASBI using electronic tools (e.g., computers and phones), referred to as electronic screening and brief intervention (e-SBI), have been developed. By reducing the amount of time required to deliver ASBI services, e-SBI could increase the uptake and reduce the costs of this service. These electronic tools may also help individuals who might otherwise be reluctant to discuss their drinking behavior with a health professional. Additionally, e-SBI can be delivered in community settings (e.g., home or school), which could increase the reach of and exposure to this intervention.

This Community Guide review focused on e-SBI interventions that are similar in structure and intent to ASBI and examined how e-SBI effectiveness varied by key intervention characteristics, in contrast to previous systematic reviews of e-SBI.^{25,26} In the review by Donoghue et al.,²⁵ the authors did not objectively define limits pertaining to the human interaction component of e-SBI in the inclusion criteria and only stratified the findings by intervention length. Moreover, the authors assessed changes in grams of ethanol per week, but did not examine e-SBI effects on other measures of alcohol use that may have greater relevance to health outcomes, such as binge drinking. Dedert and colleagues²⁶ stratified the findings by students and adults and intervention length, but not by other relevant characteristics. With different search strategies between the present review and those by Donoghue et al. and Dedert and colleagues, less than one fifth of the studies included in this review were included in the other reviews. This review is unique in that it excluded highly interactive interventions (i.e., those with more than three human-to-human sessions); those that were part of a treatment program for alcohol use disorders; those that were primarily educational programs (e.g., Alcohol Edu, College Alc); and those that only used an electronic device for monitoring drinking following traditional ASBI.

Evidence Acquisition

Community Guide methods for conducting systematic reviews and for developing evidence-based recommendations have been described elsewhere,^{27,28} and this review aligns with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) standards.²⁹ Descriptions of the conceptual approach

and methods for this systematic review, including intervention selection and outcome determination, follow.

Systematic Review Development Team

The e-SBI systematic review team (Kristin A. Tansil, Marissa B. Esser, Paramjit Sandhu, Jeffrey A. Reynolds, Randy W. Elder, Rebecca S. Williamson, Sajal K. Chattopadhyay, Michele K. Bohm, Robert D. Brewer, Lela R. McKnight-Eily, Daniel W. Hungerford, Centers for Disease Control and Prevention; Traci L. Toomey, University of Minnesota; Ralph W. Hingson, National Institute of Alcohol Abuse and Alcoholism; and Jonathan E. Fielding, Los Angeles County Department of Public Health) included representation from the Community Preventive Services Task Force (Task Force), an independent, nonfederal, unpaid panel of experts in public health and preventive medicine. Other team members included subject matter experts in ASBI, epidemiology and prevention of excessive alcohol use, and systematic review methods from the Community Guide and other government agencies and academic institutions.

Conceptual Approach and Analytic Framework

Intervention definition. To reduce excessive alcohol consumption, e-SBI uses electronic devices (e.g., computers, telephones, or mobile devices) to facilitate the delivery of key elements of traditional ASBI (as discussed in the Context section). At a minimum, e-SBI involves:

1. screening individuals for excessive drinking; and
2. delivering a brief intervention, which provides personalized feedback about the risks and consequences of excessive drinking.

Personalized feedback can be fully automated (e.g., computer-based), interactive (e.g., provided by a person via telephone), or partially automated and interactive. At least one part of the brief intervention must be delivered by an electronic device.

The brief intervention provided using e-SBI techniques may also include other common elements of traditional ASBI. One such element is motivational feedback, which has two levels: low-level motivational feedback includes general advice on how to reduce excessive alcohol consumption; high-level feedback provides more individually tailored messages based on factors such as readiness to change or developing personal goals. Another element is normative feedback, comparing an individual's own alcohol consumption with that of others (e.g., college students in the same school).

The primary aims of this review were to assess whether e-SBI reduces the prevalence, frequency, and intensity of adult binge drinking (which is reported by >90% of U.S. adult excessive drinkers),¹⁰ as well as alcohol-related harms. Secondary aims were to examine how e-SBI effectiveness varies by characteristics of the intervention, population, and setting.

Figure 1 illustrates the conceptual model developed by the review team (i.e., the authors) to indicate the causal pathways through which e-SBI, an individual-level intervention, is expected to reduce excessive alcohol consumption and related harms. The model for this review posits that implementation of e-SBI will increase the number of people screened for excessive drinking, and that those who screen

positive will receive a brief intervention. A brief intervention can lead to increased awareness of the harmful consequences of excessive drinking, which may motivate excessive drinkers to reduce their alcohol consumption, yielding a decrease in alcohol-related harms. A small proportion of those who screen positive for excessive drinking may be found to be alcohol-dependent and referred to treatment.

Inclusion criteria. To be included in the systematic review, studies had to:

1. evaluate an intervention meeting the definition of e-SBI;
2. be based on primary research published in an English-language journal, or be available as a dissertation or a technical or government report;
3. be conducted in a high-income country as defined by the World Bank³⁰;
4. include a concurrent comparison group;
5. report at least one outcome of interest, excessive alcohol consumption or alcohol-related harms; and
6. include a follow-up period > 1 month.

Exclusion criteria. Studies were excluded from this review if the interventions:

1. were not considered brief (i.e., included more than three sessions with human-to-human interactions);
2. were delivered as part of a treatment program for alcohol use disorders or as part of a broad alcohol education program;
3. involved a multifaceted health risk assessment addressing health behaviors beyond alcohol, tobacco, or other substance use; or
4. only used electronic devices to assess changes in alcohol consumption after delivering traditional ASBI.

Search for Evidence

Electronic searches for literature published from 1967 to October 2011 were conducted in the following databases: CINAHL, EconLit, Embase, ERIC, MEDLINE, NTIS, PsycINFO, Social Services Abstracts, Sociological Abstracts, Web of Science, Dissertation Abstracts, and PubMed to identify studies relevant to evaluations of e-SBI interventions. Search details are available at: www.thecommunityguide.org/alcohol/supportingmaterials/SSeSBI.html. References in all retrieved articles were examined, and additional studies identified by subject matter experts on the e-SBI systematic review team were incorporated into the review.

Assessing and Summarizing the Body of Scientific Evidence on Effectiveness

Two abstractors independently read and evaluated each study. Data from all studies that met the systematic review criteria were abstracted using an adaptation of the standardized abstraction form (original form at: www.thecommunityguide.org/about/methods.html) and coders reconciled discrepancies. Community Guide criteria for quality of study execution evaluated threats to validity, such as poor description of the intervention, study population, or sampling frame; high attrition (> 20%); or comparison groups not being comparable at baseline. Based on these criteria, studies were




Figure 1. Model showing conceptualization of how alcohol e-SBI is expected to reduce excessive alcohol consumption and related harms. e-SBI, electronic screening and brief intervention.

categorized as having good (one or no limitations), fair (two to four), or limited (four or more) quality of execution.^{27,28} Studies judged to be of limited quality of execution were excluded from the analysis.

Outcomes of Interest

Several categories of outcome measures were abstracted and summarized for the systematic review. Three binge-drinking measures, defined by the researchers in their respective studies, were assessed: prevalence (based on proportions of study participants); frequency (episodes/month); and intensity (peak alcohol consumption [maximum drinks/binge episode] or the maximum estimated blood alcohol concentration during a binge episode). Two general consumption measures were also assessed—frequency of alcohol consumption (number of drinking days/month) and total alcohol consumption (mean number of drinks/month)—as well as a measure of average drinks/drinking day. To improve comparability across studies, drinking measures were converted into standard U.S. drinks (i.e., 14 g pure alcohol/drink).

Data Analysis and Effect Estimate Calculation

Effect estimates were expressed as relative percentage changes, using the following formula:

$$[(I_{post}/C_{post}) \times (C_{pre}/I_{pre}) - 1] \times 100,$$

where I_{post} is the post-test measure for the group receiving the intervention, I_{pre} is the pretest measure for the group receiving the intervention, C_{post} is the post-test measure for the comparison/control group, and C_{pre} is the pretest measure for the comparison/control group.

If a study reported multiple follow-up measurements, then the last follow-up period up to 12 months was used to assess intervention effectiveness. If the outcomes reported in a study could not be expressed as a relative percentage, then the team qualitatively examined the evidence for direction of effect.

Effect estimates were summarized across studies using medians to reflect central tendencies and interquartile intervals to reflect variability when an adequate number of data points were available (five or more). Effect estimates were reported and summarized separately for study samples that consisted solely of people who screened positive for excessive drinking and for those that included a combination of all types of drinkers, whether or not they met criteria for excessive drinking at baseline (referred to as “all drinkers”).

Evidence Synthesis

Intervention Effectiveness

In total, 8,328 titles and abstracts were obtained (Figure 2). After removing duplicates and studies that did not meet the inclusion criteria, 37 studies that evaluated effectiveness of e-SBI for reducing excessive alcohol consumption and related harms were considered for this review. Six studies were excluded for limited quality of execution.^{31–36} The remaining 31 studies with 36 study arms were included in this review.^{37–67} All included studies were RCTs; 24 had fair quality of execution^{38,39,41,43–53,55–57,60,61,63–67} and seven had good quality.^{37,40,42,54,58,59,62} Eighteen studies had follow-up

periods of 6–12 months.^{37,39,40,43,46,47,49–55,59,60,64,65,67} Twenty-four studies (28 study arms)^{37,39,40,42,43,46–54,57–65,67} provided results for excessive drinkers only (excessive drinking criteria varied across studies) and seven studies (eight study arms)^{38,41,44,45,55,56,66} reported results for all drinkers. Details of the included studies are available at: www.thecommunityguide.org/alcohol/supplementingmaterials/SETeSBI.pdf. Analyses were conducted in 2012.

Appendix Table 1 (available online) summarizes study arm characteristics. Approximately half of the studies were conducted outside the U.S.^{37–40,43,46,47,52–54,59–61,64} and half in university settings.^{37,38,41,43–45,47,48,52,55,58,63,65,66} Nearly two thirds of the studies used probability sampling designs or attempted to recruit everyone willing to participate.^{37,39,40,43,44,46,47,52–59,62,64–67} In approximately 60% of the studies, participants were screened for excessive consumption through automated methods—most often web-based.^{38,44,47,49,52–55,58–67} More than 80% of the brief interventions were delivered solely through automated methods.^{37,38,40–45,47–56,58–64,66} Overall, 42% of the brief interventions included high-level motivational feedback (described in **Appendix Table 1**, available online).^{37,39,49–51,57–60,62,65,67} In addition, two thirds of the interventions incorporated normative feedback that compared drinking patterns among individuals with similar characteristics,^{38,40–45,47,49,50,52–55,58,59,61,63,65–67} sometimes in addition to motivational feedback. In most cases, comparison groups were assessed and given a brochure with general alcohol facts.

Summary effect estimates showed reductions in all alcohol consumption outcomes (**Table 1**). Among excessive drinkers, the largest and most consistent changes

were in frequency (median 16.5% reduction in episodes/month) and intensity of binge drinking (median 23.9% reduction in peak alcohol consumption). Peak alcohol consumption also declined in studies that included binge and non-binge drinkers in the intervention condition. Reductions in outcomes related to monthly alcohol consumption were found for studies of excessive drinkers, as well as all drinkers. Although point estimates for average number of drinks/drinking day were in the favorable direction, several studies of excessive drinkers produced nonsignificant effect estimates in the unfavorable direction.^{37,42,47,49,54} These studies reported that alcohol consumption decreased in the intervention groups; however, reductions in consumption were greater in the comparison groups. Two additional studies that assessed alcohol consumption outcomes using metrics that could not be combined with those presented in **Table 1** also showed favorable intervention effects.^{51,59}

Several studies showed that e-SBI had favorable effects on the prevalence of binge drinking and high monthly alcohol consumption. Eight studies assessed drinking prevalence,^{39,40,48,52,59,60,62,67} of which five showed a median reduction of 3.2% in binge drinking^{39,48,52,62,67} and three showed a median reduction of 12.2% in high monthly alcohol consumption.^{40,59,60} Data from two prevalence studies could not be included in the aggregated results owing to distinct reporting measures; however, both indicated that the intervention group participants were more likely to become non-excessive drinkers than the controls.^{46,64}

Fourteen studies (17 study arms)^{38,41,44,48,51–55,57,58,63,65,66} assessed the impact of e-SBI on alcohol-related harms (e.g., social, academic, and health) using self-




Figure 2. Alcohol e-SBI systematic review search yield (1967–October 2011). e-SBI, electronic screening and brief intervention.

reported alcohol problem scales. The most commonly used scale was the AUDIT,⁶⁸ which assesses quantity, frequency, and consequences of drinking. AUDIT scores range from 0 to 40, with a score ≥ 8 indicating harmful drinking. Three studies (four study arms) that assessed AUDIT scores for excessive drinkers found a median reduction of 1.1 points from a median baseline of 14.8.^{49,53,57} Similar favorable results were found for one study that assessed AUDIT score outcomes for all drinkers, finding a 0.9-point reduction from a baseline of 6.4.⁴¹ Similar results were found for other alcohol-related problem composite scales (e.g., Rutgers Alcohol Problem Index), with 12 of 17 study arms showing results in the favorable direction, seven of which were statistically significant ($p < 0.05$).^{44,51,53,54,57,65,66}

Assessment of effect modification. Table 2 shows intervention effect estimates for average drinks/drinking day stratified by descriptive variables identified by the review team as potential effect modifiers. The team selected average number of drinks/drinking day for stratified analyses because it was a commonly reported outcome measure, reported in 14 study arms.^{37–39,41,42,48,49,52–54,58,62,63} Effect estimates were in the favorable direction across all strata, with few clear differences across strata. It is noteworthy that the intervention effect

for studies with universal or probability samples appears larger than for those with convenience samples. The former provides the best estimates of the expected effects of an intervention that is brought to scale in a particular setting.

Studies with treated comparison groups. Three studies compared effectiveness of e-SBI with traditional ASBI and found mixed results.^{58,65,67} One study⁶⁷ found a 16.4% greater decrease in the proportion of people in the e-SBI group who engaged in binge drinking compared with ASBI. However, another study⁵⁸ reported a lower effectiveness of e-SBI compared with ASBI, with 15.1% more drinks consumed per drinking occasion in the e-SBI group. The third study⁶⁵ compared an e-SBI combination of Internet and face-to-face to ASBI alone, and reported a 29.5% greater reduction of average drinks per drinking occasion for e-SBI, relative to ASBI alone.

Three studies compared different forms of e-SBI feedback (e.g., personalized feedback to high-level motivational feedback), finding mixed evidence of effectiveness.^{47,64,65} Two studies^{64,65} found that motivational feedback interventions were associated with greater reductions in alcohol consumption and related problems compared with personalized feedback alone. The third study⁴⁷ showed a 13.7% reduction in mean number of

Table 1. Summary of Outcomes Related to Frequency and Quantity of Alcohol Consumption

Outcome measure	All drinkers ^a			Excessive drinkers ^b		
	Median baseline	No. of study arms	Median relative change, % (IQI)	Median baseline	No. of study arms	Median relative change, % (IQI)
Binge drinking						
Binge-drinking frequency episodes/month	3	2	-1.8	6.1	9	-16.5 (-35.6, -11.8)
Peak consumption (overall)	NA	5	-19.1 (-42.1, 17.7)	NA	9	-23.9 (-51.3, -2.1)
Peak consumption (max drinks/binge episode)	4.2	2	-33.9	9.2	5	-13.1 (-38.8, -1.8)
Peak consumption (estimated BAC)	0.10 g/dL	3	-18.1	0.18 g/dL	4	-32.2
Monthly alcohol consumption						
Frequency of alcohol consumption/month	4.6	1	-14.4	8.2	8	-11.5 (-17.3, -4.9)
Mean no. drinks/month	29.3	7	-16.2 (-33.4, -8.2)	55.6	16	-13.8 (-31.7, -10.8)
Drinks per drinking day						
Average no. of drinks/drinking day	5.1	2	-13.5	7.6	14	-5.5 (-14.5, 1.1)

^aResults for all drinkers, including those above and below threshold for excessive drinking.
^bResults for people who screened positive for excessive drinking (as defined by study authors).
 BAC, blood alcohol concentration; IQI, interquartile interval; NA, not applicable.

drinks per month, but a 2.4% increase in estimated peak blood alcohol concentration in the motivational feedback intervention compared with personalized feedback only. Because of limited evidence, further research is needed to determine whether motivational feedback interventions are associated with greater reductions in alcohol consumption.

Economic Evaluation

Evaluations of economic efficiency are conducted if the Task Force recommends an intervention. The methods and findings of the economic evaluation of alcohol e-SBI interventions to reduce excessive alcohol consumption are described in the [Appendix](#) (available online).

Conclusions

Summary of Findings

Based on the studies in this review, study participants who received e-SBI consistently reported greater reductions in excessive alcohol consumption than controls. The impact of e-SBI on excessive drinking was most pronounced for measures of binge-drinking frequency and intensity (particularly measures of peak consumption), and less pronounced for average consumption. However, some of these differences in intervention effectiveness may be due to differences in the prevalence of various drinking patterns (e.g., binge drinking versus heavy drinking), and their sensitivity for evaluating

Table 2. Average Number of Drinks per Drinking Day, Stratified by Potential Effect Modifiers

Descriptive variables	No. of study arms ^a	Median baseline	Median relative percent change in consumption (IQI)
Recruitment population			
Universal/probability sample	7 ^{39,52–54,58,62}	8.0	–8.4 (–16.8, –5.0)
Convenience	7 ^{37,38,41,42,48,49,63}	6.0	–0.8 (–16.0, 1.3)
Setting			
Health care	5 ^{39,53,54,62}	8.0	–13.0 (–23.9, –1.5)
University	8 ^{37,38,41,42,48,52,58,63}	6.7	–7.2 (–14.5, 0.6)
Community-based	1 ⁴⁹	8.8	1.3
Delivery mode of screening			
Fully automated	9 ^{38,49,52–54,58,62,63}	8.0	–6.0 (–11.5, 0.3)
Other ^b	5 ^{37,39,41,42,48}	5.7	–16.0 (–16.9, 5.8)
Delivery mode of brief intervention			
Fully automated	12 ^{37,42,48,49,52–54,58,62,63}	7.4	–6.0 (–14.5, 1.1)
Partially automated/ interpersonal ^c	1 ³⁹	3.4	–16.8
Type of feedback ^d			
Motivational			
None ^e	6 ^{48,52–54,63}	8.0	–6.0 (–16.0, –0.8)
Low-level ^f	1 ⁴²	6.0	10.6
High-level ^g	5 ^{37,39,49,58,62}	7.4	–8.4 (–23.9, –1.1)
Normative ^h	10 ^{37,42,48,49,52–54,58}	7.8	–4.2 (–9.5, 1.5)
No normative feedback	3 ^{39,62,63}	5.2	–16.8 (–31.0, –0.8)

^aIncludes only those studies for which a relative change could be calculated.

^bWritten on paper or not reported.

^cWeb-based and phone.

^dAt minimum, all types of feedback are personalized feedback.

^eNo motivational feedback—personalized risks and consequences.

^fLow-level motivational feedback (e.g., general advice).

^gHigh-level motivational feedback (e.g., commitment to change, goal setting).

^hNormative feedback—national or local comparison to others of similar age or gender; may also incorporate motivational feedback (not mutually exclusive).

IQI, interquartile interval.

changes in excessive alcohol consumption in various study populations. In studies including all drinkers, and not limited to excessive drinkers, the effectiveness of e-SBI for reducing binge drinking was less pronounced, which may partially be due to differences in the study populations. In addition, the effects of e-SBI on measures of alcohol-related harms, using measures such as Rutgers Alcohol Problem Index scores, were less pronounced; two factors may contribute to this finding. First, measurable reductions in alcohol-related problems first require reductions in alcohol consumption, and thus are less directly related to components of e-SBI. Second, categorical scales are inherently less sensitive to change than continuous measures; therefore, some changes in alcohol-related problems may have been missed by use of categorical outcomes.

The trend of the findings of this review are consistent with the findings of other systematic reviews on the effectiveness of e-SBI for reducing excessive alcohol use when measuring changes in grams or drinks of alcohol per week.^{25,26} However, the outcomes are not directly comparable because this review examined changes in the number of alcoholic drinks per month. There is little overlap in the other measures included across these e-SBI systematic reviews. For example, Dedert et al.²⁶ report no significant effects of e-SBI on binge drinking, though the measure refers to rates of binge drinking rather than changes in the frequency or quantity of binge drinking, as reported in the current study.

According to Community Guide rules of evidence,²⁷ this review found strong evidence that e-SBI is effective in reducing excessive alcohol consumption and related harms. This intervention, which focuses on individual risk reduction, can also be used in conjunction with population-level strategies to reduce excessive alcohol consumption previously recommended by the Task Force, including increasing alcohol taxes, regulating alcohol outlet density, and commercial host (dram shop) liability.⁶⁹

Limitations

This review has several limitations. First, alcohol consumption was self-reported, likely resulting in underestimation from recall bias, self-selection bias, and social desirability bias.^{37,38,41,42,45,48–51,60,61,63,70,71} Second, the validity of self-reported alcohol consumption may have been affected by the methods used to interview study participants (e.g., face-to-face, telephone, computer), and the reporting of alcohol-attributable outcomes may vary based on survey methods as well.^{72–74} Third, as is common in studies of traditional ASBI, studies in this review found substantial reductions in excessive

consumption among both intervention and control participants. This is consistent with previous studies that have found that alcohol consumption can be substantially affected by drinking assessments alone.⁵⁴ Thus, the impact of e-SBI on excessive drinking might have been even larger if intervention participants had been compared with groups that did not receive either an assessment or a brief intervention for excessive alcohol use. Fourth, exposure to e-SBI may have led participants in the intervention group to report larger changes in their alcohol consumption at follow-up than controls.⁵² However, one study⁷⁵ found larger reductions in alcohol-related injuries among e-SBI participants at the 12-month follow-up compared with controls, suggesting that e-SBI resulted in larger and sustained reductions in alcohol consumption in the treatment group.

In addition, the quality of some studies was limited because the study populations were not representative of the general population and some had high attrition rates (>20%). Attrition rates tend to be high in e-health studies,⁷⁶ including studies evaluating the effectiveness of e-SBI for reducing excessive drinking.⁷⁷ To address this problem, many studies conducted intent-to-treat analyses (i.e., applying non-completers' baseline data at follow-up), with results that were similar to participants who were retained. Lastly, this review synthesized literature that was published by October 2011, consistent with the evidence available when e-SBI was presented to the Task Force. It is possible that the magnitude of the findings would differ with the inclusion of more recent studies; however, effect estimates from more recent studies^{78,79} are consistent with those found in this review, making it unlikely that their inclusion would substantially alter the results reported in this review.

Applicability of Findings

The reviewed studies found e-SBI to be effective in most settings (e.g., healthcare settings and universities) and across various population groups. Sixteen studies (20 study arms) demonstrated favorable effects among university students, as did studies involving older populations (median age, 40 years). However, only one study included a sample of adolescents.⁶⁷ Moreover, the intervention reduced excessive alcohol consumption among men and women, although the magnitude of the sex-specific effects varied across studies.^{39,42,46,51,58,60,61} Data on the effectiveness of e-SBI by race/ethnicity and income were also limited.

Potential Harms and Other Benefits

Although e-SBI interventions pose minimal risk of harm, steps should be taken to ensure quality control and the

safeguarding of personal information. Poorly designed e-SBI applications may not incorporate validated screening measures or may provide inappropriate feedback to users, resulting in misdiagnosis of alcohol problems and delivery of ineffective advice. It is also important to assure that personal information is protected, including when e-SBI is implemented in community settings, which may not have the privacy protections required in healthcare facilities.

No additional intervention benefits were found.

Evidence Gaps

The reviewed studies demonstrated an association between e-SBI and reduced alcohol consumption and related harms; however, further research should assess e-SBI effectiveness in other settings (e.g., military, work-sites, public health organizations) and among specific populations (e.g., adolescents, racial and ethnic minority groups). Additional studies comparing e-SBI with traditional ASBI should be conducted to assess relative benefits. More research is also needed to determine the optimal intervention intensity, the relative effectiveness of different types of feedback on drinking behavior, the long-term effectiveness (beyond 12 months), and the potential usefulness of “booster sessions” to improve the long-term effect of e-SBI on drinking behavior. Future research should also evaluate the cost effectiveness of e-SBI in communities where e-SBI has been widely implemented.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

No financial disclosures were reported by the authors of this paper.

References

1. Stahre M, Roeber J, Kanny D, Brewer RD, Zhang X. Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. *Prev Chronic Dis*. 2014;11:E109. <http://dx.doi.org/10.5888/pcd11.130293>.
2. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004;291(10):1238–1245. <http://dx.doi.org/10.1001/jama.291.10.1238>.
3. Centers for Disease Control and Prevention. Alcohol-Related Disease Impact (ARDI) application. www.cdc.gov/ardi. Accessed April 19, 2016.
4. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Correction: actual causes of death in the United States, 2000. *JAMA*. 2005;293(3):293–294. <http://dx.doi.org/10.1001/jama.293.3.293>.
5. Sacks JJ, Gonzales KR, Bouchery EE, Tomedi LE, Brewer RD. 2010 National and state costs of excessive alcohol consumption. *Am J Prev Med*. 2015;49(5):e73–e79. <http://dx.doi.org/10.1016/j.amepre.2015.05.031>.
6. Centers for Disease Control and Prevention. Vital signs: binge drinking prevalence, frequency, and intensity among adults—United States, 2010. *MMWR Morb Mortal Wkly Rep*. 2012;61(1):14–19.
7. Anderson P, Cremona A, Paton A, Turner C, Wallace P. The risk of alcohol. *Addiction*. 2006;88(11):1493–1508. <http://dx.doi.org/10.1111/j.1360-0443.1993.tb03135.x>.
8. Becker U, Deis A, Sorensen TI, et al. Prediction of risk of liver disease by alcohol intake, sex, and age: a prospective population study. *Hepatology*. 1996;23(5):1025–1029. <http://dx.doi.org/10.1002/hep.510230513>.
9. Klingemann H. *Alcohol and Its Social Consequences—the Forgotten Dimension*. Geneva: WHO Regional Office for Europe; 2009. <http://dx.doi.org/10.1007/978-94-015-9725-8>.
10. Esser MB, Hedden SL, Kanny D, Brewer RD, Gfroerer JC, Naimi TS. Prevalence of alcohol dependence among U.S. adult drinkers. *Prev Chron Dis*. 2014;11:140329. <http://dx.doi.org/10.5888/pcd11.140329>.
11. Centers for Disease Control and Prevention. CDC’s alcohol screening and brief intervention efforts. www.cdc.gov/ncbddd/fasd/alcohol-screening.html. Published 2014. Accessed December 9, 2015.
12. NIH, National Institute on Alcohol Abuse and Alcoholism. *Helping Patients Who Drink Too Much: A Clinician’s Guide*. 5th ed. Bethesda, MD: U.S. DHHS; 2005.
13. McKnight-Eily LR, Liu Y, Brewer RD, et al. Vital signs: communication between health professionals and their patients about alcohol use—44 states and the District of Columbia, 2011. *MMWR Morb Morbid Wkly Rep*. 2014;63(1):16–22.
14. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption—II. *Addiction*. 1993;88(6):791–804. <http://dx.doi.org/10.1111/j.1360-0443.1993.tb02093.x>.
15. Bohn MJ, Babor TF, Kranzler HR. The Alcohol Use Disorders Identification Test (AUDIT): validation of a screening instrument for use in medical settings. *J Stud Alcohol*. 1995;56(4):423–432. <http://dx.doi.org/10.15288/jsa.1995.56.423>.
16. Smith PC, Schmidt SM, Allensworth-Davies D, Saitz R. Primary care validation of a single-question alcohol screening test. *J Gen Intern Med*. 2009;24(7):783–788. <http://dx.doi.org/10.1007/s11606-009-0928-6>.
17. Moyer VA, U.S. Preventive Services Task Force. Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2013;159(3):210–218. <http://dx.doi.org/10.7326/0003-4819-159-3-201308060-00652>.
18. Solberg LI, Maciosek MV, Edwards NM. Primary care intervention to reduce alcohol misuse ranking its health impact and cost effectiveness. *Am J Prev Med*. 2008;34(2):143–152. <http://dx.doi.org/10.1016/j.amepre.2007.09.035>.
19. U.S. DHHS: Healthfinder. Drink alcohol only in moderation. healthfinder.gov/HealthTopics/Dispatch.aspx?q1=health-conditions-and-diseases&q2=heart-health&q3=drink-alcohol-only-in-moderation. Published 2014. Accessed December 21, 2015.
20. Healthy People 2020. Substance abuse objectives. www.healthypeople.gov/2020/topicsobjectives2020/objectiveslist.aspx?topicId=40. Accessed December 21, 2015.
21. Kypri K, Saunders JB, Gallagher SJ. Acceptability of various brief intervention approaches for hazardous drinking among university students. *Alcohol Alcohol*. 2003;38(6):626–628. <http://dx.doi.org/10.1093/alcalc/agg121>.
22. Seppa K, Aalto M, Raevaara L, Perakyla A. A brief intervention for risky drinking—analysis of videotaped consultations in primary health care. *Drug Alcohol Rev*. 2004;23(2):167–170. <http://dx.doi.org/10.1080/09595230410001704145>.
23. Hingson RW, Heeren T, Edwards EM, Saitz R. Young adults at risk for excess alcohol consumption are often not asked or counseled about

- drinking alcohol. *J Gen Intern Med.* 2012;27(2):179–184. <http://dx.doi.org/10.1007/s11606-011-1851-1>.
24. Yoast RA, Wilford BB, Hayashi SW. Encouraging physicians to screen for and intervene in substance use disorders: obstacles and strategies for change. *J Addict Dis.* 2008;27(3):77–97. <http://dx.doi.org/10.1080/10550880802122687>.
 25. Donoghue K, Patton R, Phillips T, Deluca P, Drummond C. The effectiveness of electronic screening and brief intervention for reducing levels of alcohol consumption: a systematic review and meta-analysis. *J Med Internet Res.* 2014;16(6):e142. <http://dx.doi.org/10.2196/jmir.3193>.
 26. Dedert EA, McDuffie JR, Stein R, et al. Electronic interventions for alcohol misuse and alcohol use disorders: a systematic review. *Ann Intern Med.* 2015;163(3):205–2214. <http://dx.doi.org/10.7326/M15-0285>.
 27. Briss PA, Zaza S, Pappaioanou M, et al. Developing an evidence-based Guide to Community Preventive Services—methods. *Am J Prev Med.* 2000;18(1 Suppl):35–43. [http://dx.doi.org/10.1016/S0749-3797\(99\)00119-1](http://dx.doi.org/10.1016/S0749-3797(99)00119-1).
 28. Zaza S, Wright-De Aguero LK, Briss PA, et al. Data collection instrument and procedure for systematic reviews in the Guide to Community Preventive Services. *Am J Prev Med.* 2000;18(1 Suppl):44–74. [http://dx.doi.org/10.1016/S0749-3797\(99\)00122-1](http://dx.doi.org/10.1016/S0749-3797(99)00122-1).
 29. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med.* 2009;151(4):264–269, W264.
 30. World Bank. High-income economies. http://data.worldbank.org/about/country-classifications/country-and-lending-groupsHigh_income. Published 2014. Accessed December 21, 2015.
 31. Doumas DM, Anderson J. Reducing alcohol use in first-year university students: evaluation of a web-based personalized feedback program. *J Coll Counsel.* 2009;12(1):18–32. <http://dx.doi.org/10.1002/j.2161-1882.2009.tb00037.x>.
 32. Doumas DM, Haustveit T. Reducing heavy drinking in intercollegiate athletes: evaluation of a web-based personalized feedback program. *Sport Psychol.* 2008;22:212–228.
 33. Rose GL, MacLean CD, Skelly J, Badger GJ, Ferraro TA, Helzer JE. Interactive voice response technology can deliver alcohol screening and brief intervention in primary care. *J Gen Intern Med.* 2010;25(4):340–344. <http://dx.doi.org/10.1007/s11606-009-1233-0>.
 34. Saitz R, Palfai TP, Freedner N, et al. Screening and brief intervention online for college students: the ihealth study. *Alcohol Alcohol.* 2007;42(1):28–36. <http://dx.doi.org/10.1093/alcac/agl092>.
 35. Weitzel JA, Bernhardt JM, Usdan S, Mays D, Glanz K. Using wireless handheld computers and tailored text messaging to reduce negative consequences of drinking alcohol. *J Stud Alcohol Drugs.* 2007;68(4):534–537. <http://dx.doi.org/10.15288/jasad.2007.68.534>.
 36. Williams J, Herman-Stahl M, Calvin SL, Pemberton M, Bradshaw M. Mediating mechanisms of a military web-based alcohol intervention. *Drug Alcohol Depend.* 2009;100(3):248–257. <http://dx.doi.org/10.1016/j.drugaldep.2008.10.007>.
 37. Bewick BM, West R, Gill J, et al. Providing web-based feedback and social norms information to reduce student alcohol intake: a multisite investigation. *J Med Internet Res.* 2010;12(5):e59. <http://dx.doi.org/10.2196/jmir.1461>.
 38. Bewick BM, Trusler K, Mulhern B, Barkham M, Hill AJ. The feasibility and effectiveness of a web-based personalised feedback and social norms alcohol intervention in UK university students: a randomised control trial. *Addict Behav.* 2008;33(9):1192–1198. <http://dx.doi.org/10.1016/j.addbeh.2008.05.002>.
 39. Bischof G, Grothues JM, Reinhardt S, Meyer C, John U, Rumpf HJ. Evaluation of a telephone-based stepped care intervention for alcohol-related disorders: a randomized controlled trial. *Drug Alcohol Depend.* 2008;93(3):244–251. <http://dx.doi.org/10.1016/j.drugaldep.2007.10.003>.
 40. Boon B, Risselada A, Huiberts A, Riper H, Smit F. Curbing alcohol use in male adults through computer generated personalized advice: randomized controlled trial. *J Med Internet Res.* 2011;13(2):e43. <http://dx.doi.org/10.2196/jmir.1695>.
 41. Bryant ZE. *Testing the Effectiveness of E-Mailed BASICS Feedback with College Students [dissertation]*. Auburn: Auburn University; 2009.
 42. Chiauzzi E, Green TC, Lord S, Thum C, Goldstein M. My student body: a high-risk drinking prevention web site for college students. *J Am Coll Health.* 2005;53(6):263–274. <http://dx.doi.org/10.3200/JACH.53.6.263-274>.
 43. Cunningham JA, Wild TC, Cordingley J, Van Mierlo T, Humphreys K. Twelve-month follow-up results from a randomized controlled trial of a brief personalized feedback intervention for problem drinkers. *Alcohol Alcohol.* 2010;45(3):258–262. <http://dx.doi.org/10.1093/alcac/agq009>.
 44. Doumas DM, Kane CM, Navarro TB, Roman J. Decreasing heavy drinking in first-year students: evaluation of a web-based personalized feedback program administered during orientation. *J Coll Counsel.* 2011;14(1):5–20. <http://dx.doi.org/10.1002/j.2161-1882.2011.tb00060.x>.
 45. Doumas DM, Haustveit T, Coll KM. Reducing heavy drinking among first year intercollegiate athletes: a randomized controlled trial of web-based normative feedback. *J Appl Sport Psychol.* 2010;22(3):247–261. <http://dx.doi.org/10.1080/10413201003666454>.
 46. Eberhard S, Nordstrom G, Hoglund P, Ojehagen A. Secondary prevention of hazardous alcohol consumption in psychiatric out-patients: a randomised controlled study. *Soc Psychiatry Psychiatr Epidemiol.* 2009;44(12):1013–1021. <http://dx.doi.org/10.1007/s00127-009-0023-7>.
 47. Ekman DS, Andersson A, Nilsson P, Stahlbrandt H, Johansson AL, Bendtsen P. Electronic screening and brief intervention for risky drinking in Swedish university students—a randomized controlled trial. *Addict Behav.* 2011;36(6):654–659. <http://dx.doi.org/10.1016/j.addbeh.2011.01.015>.
 48. Hedman AS. *Effects of Personalized Feedback and Tailored Health Communication on Alcohol Consumption, Alcohol-Related Behaviors, and Attitude Among Binge Drinking College Students [dissertation]*. Lawrence: Department of Health, Sports, and Exercise Science, University of Kansas; 2007.
 49. Hester RK, Squires DD, Delaney HD. The drinker's check-up: 12-month outcomes of a controlled clinical trial of a stand-alone software program for problem drinkers. *J Subst Abuse Treat.* 2005;28(2):159–169. <http://dx.doi.org/10.1016/j.jsat.2004.12.002>.
 50. Hester RK, Delaney HD. Behavioral self-control program for Windows: results of a controlled clinical trial. *J Consult Clin Psychol.* 1997;65(4):686–693. <http://dx.doi.org/10.1037/0022-006X.65.4.686>.
 51. Hester RK, Delaney HD, Campbell W. ModerateDrinking.Com and moderation management: outcomes of a randomized clinical trial with non-dependent problem drinkers. *J Consult Clin Psychol.* 2011;79(2):215–224. <http://dx.doi.org/10.1037/a0022487>.
 52. Kypri K, Hallett J, Howat P, et al. Randomized controlled trial of proactive web-based alcohol screening and brief intervention for university students. *Arch Intern Med.* 2009;169(16):1508–1514. <http://dx.doi.org/10.1001/archinternmed.2009.249>.
 53. Kypri K, Langley JD, Saunders JB, Cashell-Smith ML, Herbison P. Randomized controlled trial of web-based alcohol screening and brief intervention in primary care. *Arch Intern Med.* 2008;168(5):530–536. <http://dx.doi.org/10.1001/archinternmed.2007.109>.
 54. Kypri K, Saunders JB, Williams SM, et al. Web-based screening and brief intervention for hazardous drinking: a double-blind randomized controlled trial. *Addiction.* 2004;99(11):1410–1417. <http://dx.doi.org/10.1111/j.1360-0443.2004.00847.x>.
 55. Martens MP, Kilmer JR, Beck NC, Zamboanga BL. The efficacy of a targeted personalized drinking feedback intervention among intercollegiate athletes: a randomized controlled trial. *Psychol Addict Behav.* 2010;24(4):660–669. <http://dx.doi.org/10.1037/a0020299>.

56. Matano RA, Koopman C, Wanat SF, et al. A pilot study of an interactive web site in the workplace for reducing alcohol consumption. *J Subst Abuse Treat*. 2007;32(1):71–80. <http://dx.doi.org/10.1016/j.jsat.2006.05.020>.
57. Mello MJ, Longabaugh R, Baird J, Nirenberg T, Woolard R. DIAL: a telephone brief intervention for high-risk alcohol use with injured emergency department patients. *Ann Emerg Med*. 2008;51(6):755–764. <http://dx.doi.org/10.1016/j.annemergmed.2007.11.034>.
58. Mignogna J. A test of moderating factors of brief interventions for hazardous alcohol use among college students. *DAI: Sci Engineer*. 2011;71(10-B):6446.
59. Neumann T, Neuner B, Weiss-Gerlach E, et al. The effect of computerized tailored brief advice on at-risk drinking in subcritically injured trauma patients. *J Trauma*. 2006;61(4):805–814. <http://dx.doi.org/10.1097/01.ta.0000196399.29893.52>.
60. Riper H, Kramer J, Smit F, Conijn B, Schippers G, Cuijpers P. Web-based self-help for problem drinkers: a pragmatic randomized trial. *Addiction*. 2008;103(2):218–227. <http://dx.doi.org/10.1111/j.1360-0443.2007.02063.x>.
61. Spijkerman R, Roek MA, Vermulst A, Lemmers L, Huiberts A, Engels RC. Effectiveness of a web-based brief alcohol intervention and added value of normative feedback in reducing underage drinking: a randomized controlled trial. *J Med Internet Res*. 2010;12(5):e65. <http://dx.doi.org/10.2196/jmir.1465>.
62. Suffoletto B, Callaway C, Kristan J, Kraemer K, Clark DB. Text-message-based drinking assessments and brief interventions for young adults discharged from the emergency department. *Alcohol Clin Exp Res*. 2011;36(3):552–560. <http://dx.doi.org/10.1111/j.1530-0277.2011.01646.x>.
63. Sugarman DE. Web-based alcohol feedback intervention for heavy drinking college students: does drinking control strategy use mediate intervention effects? *DAI: Sci Engineer*. 2010;70(11-B):7222.
64. Trinks A, Festin K, Bendtsen P, Nilsen P. Reach and effectiveness of a computer-based alcohol intervention in a Swedish emergency room. *Int Emerg Nurs*. 2010;18(3):138–146. <http://dx.doi.org/10.1016/j.ienj.2009.08.004>.
65. Walters ST, Vader AM, Harris TR, Field CA, Jouriles EN. Dismantling motivational interviewing and feedback for college drinkers: a randomized clinical trial. *J Consult Clin Psychol*. 2009;77(1):64–73. <http://dx.doi.org/10.1037/a0014472>.
66. Walters ST, Vader AM, Harris TR. A controlled trial of web-based feedback for heavy drinking college students. *Prev Sci*. 2007;8(1):83–88. <http://dx.doi.org/10.1007/s11121-006-0059-9>.
67. Walton MA, Chermack ST, Shope JT, et al. Effects of a brief intervention for reducing violence and alcohol misuse among adolescents: a randomized controlled trial. *JAMA*. 2010;304(5):527–535. <http://dx.doi.org/10.1001/jama.2010.1066>.
68. Babor TF, De la Fuente JR, Saunders J, Grant M. *Audit. The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Care*, 2nd ed. Geneva, Switzerland: WHO; 1992.
69. Guide to Community Preventive Services. Preventing excessive alcohol consumption. www.thecommunityguide.org/alcohol/index.html. Published 2015. Accessed December 21, 2015.
70. Nelson DE, Naimi TS, Brewer RD, Roeder JUS. State alcohol sales compared to survey data, 1993–2006. *Addiction*. 2010;105(9):1589–1596. <http://dx.doi.org/10.1111/j.1360-0443.2010.03007.x>.
71. Babor TF, Steinberg K, Anton R, Del Boca F. Talk is cheap: measuring drinking outcomes in clinical trials. *J Stud Alcohol*. 2000;61(1):55–63. <http://dx.doi.org/10.15288/jsa.2000.61.55>.
72. Midanik LT, Greenfield TK. Interactive voice response versus computer-assisted telephone interviewing (CATI) surveys and sensitive questions: the 2005 National Alcohol Survey. *J Stud Alcohol Drugs*. 2008;69(4):580–588. <http://dx.doi.org/10.15288/jsad.2008.69.580>.
73. Midanik LT, Greenfield TK, Rogers JD. Reports of alcohol-related harm: telephone versus face-to-face interviews. *J Stud Alcohol*. 2001;62(1):74–78. <http://dx.doi.org/10.15288/jsa.2001.62.74>.
74. Sinadinovic K, Wennberg P, Berman AH. Population screening of risky alcohol and drug use via Internet and Interactive Voice Response (IVR): a feasibility and psychometric study in a random sample. *Drug Alcohol Depend*. 2011;114(1):55–60. <http://dx.doi.org/10.1016/j.drugalcdep.2010.09.004>.
75. Mello MJ, Baird J, Nirenberg TD, Lee C, Woolard R, Longabaugh R. DIAL: a randomised trial of a telephone brief intervention for alcohol. *Inj Prev*. 2013;19(1):44–48. <http://dx.doi.org/10.1136/injuryprev-2012-040334>.
76. Eysenbach G. The law of attrition. *J Med Internet Res*. 2005;7(1):e11. <http://dx.doi.org/10.2196/jmir.7.1.e11>.
77. Postel MG, de Haan HA, ter Huurne ED, van der Palen J, Becker ES, de Jong CA. Attrition in web-based treatment for problem drinkers. *J Med Internet Res*. 2011;13(4):e117. <http://dx.doi.org/10.2196/jmir.1811>.
78. Kypri K, Vater T, Bowe SJ, et al. Web-based alcohol intervention for Māori university students. *Addiction*. 2013;108(2):331–338. <http://dx.doi.org/10.1111/j.1360-0443.2012.04067.x>.
79. Kypri K, Vater T, Bowe SJ, et al. Web-based alcohol screening and brief intervention for university students: a randomized trial. *JAMA*. 2014;311(12):1218–1224. <http://dx.doi.org/10.1001/jama.2014.2138>.

Appendix

Supplementary data

Supplementary data associated with this article can be found at <https://www.thecommunityguide.org/sites/default/files/assets/eac-ajpm-app-eSBI.pdf>.